


Anti-Human CD196 (CCR6) PE

Catalog Number: 12-1969

Also Known As:

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD19 APC (cat. 17-0199) and Mouse IgG1 kappa Isotype Control PE (cat. 12-4714) (left) or Anti-Human CD196 (CCR6) PE (right). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Human CD196 (CCR6) PE


 Catalog Number: 12-1969

Clone: R6H1


Concentration: 5 µL (0.06 µg)/test

Host/Isotype: Mouse IgG1

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

 Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

This R6H1 monoclonal antibody reacts with CD196 (also known as CCR6), a seven transmembrane G protein-coupled receptor expressed on T, B, dendritic, natural killer, and Langerhans cells. This CC chemokine receptor uniquely binds MIP-3a/CCL20, a chemoattractant for dendritic cells, effector/memory T cells, and B cells. CD196 is also involved in host defense and inflammation at epithelial sites. Furthermore, this receptor has been implicated in Th17 differentiation and CD4+FoxP3+ regulatory T cell development.

Applications Reported

This R6H1 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This R6H1 antibody has been pre-titrated and tested by flow cytometric analysis on normal human peripheral blood cells. This can be used at 5 µL (0.06 µg)/per test. A test is defined as the amount (µg)/test of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

References

Liu H, Rohowsky-Kochan C. Regulation of IL-17 in human CCR6+ effector memory T cells. *J Immunol.* 2008 Jun 15;180(12):7948-57

Singh SP, Zhang HH, Foley JF, Hedrick MN, Farber JM. Human T cells that are able to produce IL-17 express the chemokine receptor CCR6. *J Immunol.* 2008 Jan 1;180(1):214-21

Lim HW, Lee J, Hillsamer P, Kim CH. Human Th17 cells share major trafficking receptors with both polarized effector T cells and FOXP3+ regulatory T cells. *J Immunol.* 2008 Jan 1;180(1):122-9

Shutyser E, Struyf S, Van Damme J. The CC chemokine CCL20 and its receptor CCR6. *Cytokine Growth Factor Rev.* 2003 Oct;14(5):409-26.

Carramolino L, Kremer L, Goya I, Varona R, Buesa JM, Gutiérrez J, Zaballo A, Martínez-A C, Márquez G. Down-regulation of the beta-chemokine receptor CCR6 in dendritic cells mediated by TNF-alpha and IL-4. *J Leukoc Biol.* 1999 Nov;66(5):837-44. (clone R6H1, FC, Pubmed)

Related Products

12-4714 Mouse IgG1 K Isotype Control PE (P3.6.2.8.1)

17-0199 Anti-Human CD19 APC (HIB19)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com