


Anti-Human CD197 (CCR7) PE

Catalog Number: 12-1979

Also known as: EBI-1, MIP-3 beta Receptor

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD45RA APC (cat. 17-0458) and Rat IgG2a K Isotype Control PE (cat. 12-4321) (left) or Anti-Human CD197 (CCR7) PE (right). Cells in the lymphocyte gate were used for analysis.

Product Information



Contents: Anti-Human CD197 (CCR7) PE

Catalog Number: 12-1979

Clone: 3D12

Concentration: 5 μ L (0.125 μ g)/test

Host/Isotype: Rat IgG2a, kappa



LOT



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Caution, contains Azide

Description

The 3D12 monoclonal antibody reacts with human CCR7, also known as EBI-1 and CD197. CCR7 is a member of the G-protein-coupled chemokine receptor family with seven membrane-spanning domains and functions as a receptor for 6CKine/SLC (secondary lymphoid-tissue chemokine), CCL19 and CCL21. CCR7 has been shown to be internalized via clathrin-coated pits and the majority recycled back to the plasma membrane. CCR7 is expressed on T cells and can be used to distinguish populations of naïve from central and effector memory T cells. CCR7 has been shown to play a role in migration of memory T cells to inflamed tissue. Expression of CCR7 is also found on DC's. During DC maturation CCR7 expression increases and is thought to be involved in a variety of functions: chemotaxis to the lymph node, cellular architecture, rate of endocytosis, survival and maturation. Expression of CCR7 on the cell surface can be down regulated upon ligand binding.

Applications Reported

This 3D12 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This 3D12 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 μ L (0.125 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

It is recommended that the staining incubation time be increased to at least 45 minutes at 4°C for optimal staining.

References

Geginat J, Lanzavecchia A, Sallusto F. Proliferation and differentiation potential of human CD8+ memory T-cell subsets in response to antigen or homeostatic cytokines. *Blood*. 2003 Jun 1;101(11):4260-6. (3D12, FC, PubMed)

Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature*. 1999 Oct 14;401(6754):708-12. (3D12, FC, PubMed)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Human CD197 (CCR7) PE

Catalog Number: 12-1979

Also known as: EBI-1, MIP-3 beta Receptor

RUO: For Research Use Only. Not for use in diagnostic procedures.

Related Products

12-1971 Anti-Mouse CD197 (CCR7) PE (4B12)

12-4321 Rat IgG2a K Isotype Control PE (eBR2a)

17-0458 Anti-Human CD45RA APC (HI100)