


Anti-Human/Mouse beta-Catenin PE

Catalog Number: 12-2567

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of Jurkat cell line with Mouse IgG1 K Isotype Control PE (cat. 12-4714) (blue histogram) or Anti-Human/Mouse beta-Catenin PE (purple histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Human/Mouse beta-Catenin PE

REF **Catalog Number:** 12-2567

Clone: 15B8

Concentration: 5 μ L (0.125 μ g)/test

Host/Isotype: Mouse IgG1, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial



Description

The 15B8 monoclonal antibody reacts with human and mouse beta-catenin, one member of a family of catenins, which are intracellular proteins that interact with cadherins to mediate cellular adhesion. More specifically, beta-catenin binds to the cytoplasmic tail of E-cadherin. In addition, this molecule is a component of the canonical Wnt signaling pathway. In the absence of Wnt binding its receptor, beta-catenin is phosphorylated and resides in the cytoplasm where it is eventually targeted for degradation by ubiquitination. Upon Wnt binding, beta-catenin becomes dephosphorylated, translocates to the nucleus, and modulates gene expression in partnership with the transcription factors T cell factor (TCF) and lymphocyte enhancer binding factor (LEF). Expression of beta-catenin is found in a wide variety of non-immune and immune tissues, including thymocytes and T and B lymphocytes. The Wnt & beta-catenin signaling pathway has been demonstrated to play a crucial role in the development of T, B, and hematopoietic stem cells.

Applications Reported

This 15B8 antibody has been reported for use in intracellular staining followed by flow cytometric analysis.

Applications Tested

This 15B8 antibody has been pre-titrated and tested by intracellular staining and flow cytometric analysis of Jurkat cell line using the Foxp3 Staining Buffer Set (cat. 00-5523). This can be used at 5 μ L (0.125 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Xu M, Sharma A, Hossain MZ, Wiest DL, Sen JM. Sustained expression of pre-TCR induced beta-catenin in post-beta-selection thymocytes blocks T cell development. *J Immunol.* 2009 Jan 15;182(2):759-65.

Staal FJ, Sen JM. The canonical Wnt signaling pathway plays an important role in lymphopoiesis and hematopoiesis. *Eur J Immunol.* 2008 Jul;38(7):1788-94. Review.

Xu, Y, Banerjee D, Huelsken J, Birchmeier W, and Sen JM. Deletion of beta-catenin impairs T cell development. *Nat.*

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Human/Mouse beta-Catenin PE

Catalog Number: 12-2567

RUO: For Research Use Only. Not for use in diagnostic procedures.

Immunol. 2003 4:1177-1182.

Hoschuetzky H, Aberle H, Kemler R. Beta-catenin mediates the interaction of the cadherin-catenin complex with epidermal growth factor receptor. J Cell Biol. 1994 Dec;127(5):1375-80.

Related Products

00-5523 Foxp3 / Transcription Factor Staining Buffer Set

12-4714 Mouse IgG1 K Isotype Control PE (P3.6.2.8.1)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com