


Anti-Human CD85g (ILT7) PE

Catalog Number: 12-5179

Also known as: ILT-7, LILRA4

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD123 APC (cat. 17-1239) (left) or Anti-Human CD303a FITC (cat. 11-9818) (right) and Anti-Human CD85g (ILT7) PE (right). Quadrant lines demarcate isotype control (cat. 12-4714).

Product Information



Contents: Anti-Human CD85g (ILT7) PE

Catalog Number: 12-5179

Clone: eBio17G10.2 (17G10.2)

Concentration: 5 μ L (0.125 μ g)/test

Host/Isotype: Mouse IgG1, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

Batch Code: Refer to vial



Use By: Refer to vial

Contains sodium azide

Description

The eBio17G10.2 monoclonal antibody reacts with human immunoglobulin-like transcript 7 (ILT7), also known as CD85g. ILT7 is part of the leukocyte immunoglobulin-like receptor (LIR) gene family and associates with Fc epsilon R1 gamma to form a functional complex which transduces ITAM-mediated signals that negatively modulates TLR-induced type I IFN production by human plasmacytoid dendritic cells. ILT7 is expressed exclusively on plasmacytoid dendritic cells and is absent in monocyte derived DCs. Treatment with IL-3 or CpG causes a decrease in expression.

Double staining of human PBMCs with eBio17G10.2 and anti-pDC marker BDCA2 yields virtually complete co-staining. Cross linking of ILT7 with eBio17G10.2 antibody causes reduced transcription of IFN alpha1, IFN alpha2, IFN alpha4 and IFN beta.

Applications Reported

This eBio17G10.2 (17G10.2) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBio17G10.2 (17G10.2) antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 μ L (0.125 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Cao, W., et al. 2006. Plasmacytoid dendritic cell-specific receptor ILT7-FcRI inhibits Toll-like receptor-induced interferon production. *J. Exp. Med.* 203: 1399-1405.

Ju XS, Hacker C, Scherer B, Redecke V, Berger T, Schuler G, Wagner H, Lipford GB, Zenke M. Immunoglobulin-like transcripts ILT2, ILT3 and ILT7 are expressed by human dendritic cells and down-regulated following activation. *Gene.* 2004 Apr 28;331:159-64.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Human CD85g (ILT7) PE

Catalog Number: 12-5179

Also known as: ILT-7, LILRA4

RUO: For Research Use Only. Not for use in diagnostic procedures.

Rissoan MC, Duhon T, Bridon JM, Bendriss-Vermare N, Peronne C, de Saint Vis B, Briere F, Bates EE. Subtractive hybridization reveals the expression of immunoglobulin-like transcript 7, Eph-B1, granzyme B, and 3 novel transcripts in human plasmacytoid dendritic cells. *Blood*. 2002 Nov 1;100(9):3295-303.

Related Products

11-9818 Anti-Human CD303a FITC (201A)

12-4714 Mouse IgG1 K Isotype Control PE (P3.6.2.8.1)

17-1239 Anti-Human CD123 APC (6H6)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com