

Alexa Fluor® 647 anti-mouse CD55 (DAF)

Catalog # / Size: 131805 / 25 µg
131806 / 100 µg

Clone: RIKO-3

Isotype: Armenian Hamster IgG

Immunogen: Transfected cells


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions. The solution is free of unconjugated Alexa Fluor® 647.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 splenocytes stained with RA3-6B2 (B220) PE and Riko-3 Alexa Fluor® 647

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10^6 cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

** Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

Description: CD55, also known as DAF (decay accelerating factor), is expressed on the surfaces of red blood cells, hematopoietic cells and many non-hematopoietic cells. There are two isoforms of mouse CD55, a GPI-anchored isoform and a transmembrane isoform. Both are recognized by RIKO-3. The function of CD55 is to protect damage by complement deposition.

Antigen References: 1. Miwa T *et al.* 2001. *Immunology* 104:207
2. Lin FY *et al.* 2001 *Immunology* 104:215

Related Products:Product

Alexa Fluor® 647 Armenian Hamster IgG Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
TruStain fcX™ (anti-mouse CD16/32)

Clone
HTK888

93

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.