

PE anti-mouse CD66a (CEACAM1a)

Catalog # / Size: 134505 / 25 µg
134506 / 100 µg

Clone: MAb-CC1

Isotype: Mouse IgG1

Immunogen: Balb/c mouse purified intestinal brush border membrane


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Balb/c mouse splenocytes stained with MAb-CC1 PE (lower panel) or mouse IgG1 PE isotype control (upper panel) and CD45R/B220 (RA3-6B2) APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

1. Turner BC, *et al.* 2004. *J. Virol.* 78 (10):5486
2. Williams RK, *et al.* 1990. *J. Virol.* 64:3817
3. Dveksler GS, *et al.* 1993. *Proc. Natl. Acad. Sci. USA.* 90:1716


Description: CD66a, known as CEACAM1a, carcinoembryonic antigen-related cell adhesion molecule 1a, is a glycoprotein of the immunoglobulin superfamily and the carcinoembryonic antigen family. Isoforms expressing either two or four alternatively spliced Ig-like domains in mice have been found in a number of epithelial, endothelial, or hematopoietic tissues. CEACAM1a functions as an intercellular adhesion molecule, an angiogenic factor, and a tumor cell growth inhibitor. It also serves as a signal regulatory protein influencing B cell receptor complex-mediated activation. Moreover, the mouse CEACAM1a protein is targets of viral or bacterial pathogens. It was reported that targeted disruption of the CEACAM1a gene resulting in a partial ablation of the protein in mice led to reduced susceptibility to virus infection. The antibody recognizes the N-terminal domain of murine CEACAM1a, does not recognize murine CEACAM1b, an allele in SJL mice.

Antigen References:

1. Nakagaki K, *et al.* 2005. *J. Virol.* 79(10):6102
2. Greicius G *et al.* 2003. *J. Leukoc. Biol.* 74(1):126
3. Hemmila E *et al.* 2004. *J. Virol.* 78(18):10156

Related Products:

Product	Clone	Application
PE Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.