

APC anti-mouse CD68

Catalog # / Size: 137007 / 25 µg
137008 / 100 µg

Clone: FA-11

Isotype: Rat IgG2a

Immunogen: Purified Con A receptor glycoproteins from the P815 cell line


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Thioglycolate-elicited Balb/c peritoneal macrophages intracellularly stained with FA-11 APC

Applications:

Applications: FC, ICFC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.06 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported (for relevant formats) applications include: immunoprecipitation^{1, 2}, Western Blot^{1, 2}, and immunohistochemical staining of frozen section².

Application References: 1. Silva RP, *et al.* 1999. *Biochem. J.* 338:687. (IP WB)
2. Rabinowitz SS, *et al.* 1991. *J. Exp. Med.* 174:827. (IP WB IHC)

Description: Mouse CD68, also known as macrosialin, is an 85-115 kD member of the lysosomal-associated membrane protein (LAMP) family. It is a heavily glycosylated and predominantly intracellular protein, mainly in late endosomes. Macrosialin is the murine homolog to the human macrophage glycoprotein CD68. It is expressed on tissue macrophages, Langerhans cells and at low levels on dendritic cells. Lamp proteins may have functions relating to cell-cell interaction or cell-ligand interaction. The biological function of CD68 is not completely understood.

Antigen References: 1. Ramprasad MP, *et al.* 1996. *Proc. Natl. Acad. Sci. USA* 93:14833.
2. Smith MJ, *et al.* 1987. *J. Cell. Sci.* 87:113.

Related Products:	Product	Clone	Application
	APC Rat IgG2a, κ Isotype Ctrl	RTK2758	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.