


Anti-Mouse Dendritic Cell Marker (33D1) PE

Catalog Number: 12-5884

Also Known As: DC Marker, DC inhibitor receptor 2

RUO: For Research Use Only



Staining of BALB/c splenocytes with Anti-Mouse MHC Class II (I-A/I-E) FITC (cat. 11-5321) and 0.125 ug of Rat IgG2b K Isotype Control PE (cat. 12-4032) (left) or 0.125 ug of Anti-Mouse Dendritic Cell Marker (33D1) PE (right). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse Dendritic Cell Marker (33D1) PE


REF **Catalog Number:** 12-5884

Clone: 33D1

Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2b, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

 **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

Description

The 33D1 monoclonal antibody reacts with a mouse DC-specific surface marker. The nature and biological activity of the 33D1 antigen is as of yet unknown. 33D1 has been reported on a variety of dendritic cell subpopulations from mouse thymus, spleen, lymph node, and Peyer's patch. Bone marrow dendritic cells require GM-CSF to express the 33D1 antigen and this expression is down regulated in the presence of IL-4. 33D1 antigen has been detected *in vivo* in brain dendritic cells post infection via *Toxoplasma gondii*.

Applications Reported

The 33D1 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The 33D1 antibody has been tested by flow cytometric analysis of mouse splenocyte cell suspensions. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Itoyaga J, Lubkin A, Fiorese C, Lahoud MH, Caminschi I, Huang Y, Rodriguez A, Clausen BE, Park CG, Trumfheller C, Steinman RM. Comparable T helper 1 (Th1) and CD8 T-cell immunity by targeting HIV gag p24 to CD8 dendritic cells within antibodies to Langerin, DEC205, and Clec9A. *Proc Natl Acad Sci U S A*. 2011 Feb 8;108(6):2384-9. (DCIR2, **33D1**, FC, PubMed)

Dudziak D, Kamphorst AO, Heidkamp GF, Buchholz VR, Trumfheller C, Yamazaki S, Cheong C, Liu K, Lee HW, Park CG, Steinman RM, Nussenzweig MC. Differential antigen processing by dendritic cell subsets in vivo. *Science*. 2007 Jan 5;315(5808):107-11. (**33D1**, IHC frozen, PubMed)

Steinman, R.M., B. Gutchinov, M.D. Witmer, and M.C. Nussenzweig. 1983. Dendritic cells are the principal stimulators of the primary mixed leukocyte reaction in mice. *J. Exp. Med.* 157: 613 - 627.

Nussenzweig, M.C., R.M. Steinman, M.D. Witmer, and B. Gutchinov. 1982. A monoclonal antibody specific for mouse dendritic cells. *Proc. Natl. Acad. Sci. USA* 79: 161 - 165.

Related Products

11-0031 Anti-Mouse CD3e FITC (145-2C11)

11-5321 Anti-Mouse MHC Class II (I-A/I-E) FITC (M5/114.15.2)

12-4031 Rat IgG2b K Isotype Control PE
17-2051 Anti-Mouse CD205 APC (205yekta)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com