

## **Product Data Sheet**

## PE anti-mouse CD73

Catalog # / Size:	127205 / 25 μg 127206 / 100 μg			
Clone:	TY/11.8			x 600
Isotype:	Rat IgG1, κ		1.4 4 1	
Immunogen:	CHO cells transfected with mouse CD73	ω.		
Reactivity:	Mouse	1.8 PE		Alexan
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.	TY/11.8		
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.			Sec. 1
Concentration:	0.2 mg/ml	-		
Storage:	The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. <b>Do not freeze.</b>	CD3 (145-2C11) APC BALB/c mouse splenocytes stained with TY/11.8 PE and CD3 (145-2C11) APC		
Application	IS:			
Applications:	FC - Quality tested			
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining immunofluorescent staining, the suggested use of this reagent is $\leq 0.25 \ \mu g \ p c$ recommended that the reagent be titrated for optimal performance for each approximate the suggested statement of the sugge	er 10 <sup>6</sup>	cells in 100	etric analysis. For μl volume. It is
Application References:	1. Yamshita Y, et al 1998. Eur. J. Immunol. 28:2981.			
Description:	CD73 (ecto-5`-nucleotidase) is a 69 kD GPI-anchored surface protein. In mice restricted to CD11b+ myeloid cells. In spleen, it is largely expressed on T cell		ression of C	D73 in bone marrow
Antigen References:	1 Yamashita Y et al 1998 Fur J Immunol 28:2981			

Antigen References: 1. Yamashita Y, et al 1998. Eur. J. Immunol. 28:2981. 2. Resta R, et al. 1998. Immunol. Rev. 161:95.

## **Related Products: Product**

PE Rat IgG1, κ Isotype Ctrl Cell Staining Buffer RBC Lysis Buffer (10X) TruStain fcX™ (anti-mouse CD16/32) Clone RTK2071

93

Application FC, ICFC FC, ICC, ICFC FC, ICFC FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.