

Purified anti-mouse CD88 (C5aR)

Catalog # / Size: 135802 / 100 µg

Clone: 20/70

Isotype: Rat IgG2b, κ

Immunogen: mC5aR transfected cells


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



C57BL/6 bone marrow cells stained with 20/70 PE and Ly-6G/Ly-6C (Gr-1) APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications for the relevant formats include: blocking of C5a binding.

Application References:

1. Godau J, *et al.* 2004. *J. Immunol.* 173:3437.
2. Shushakova N, *et al.* 2002. *J. Clin. Invest.* 110:1823.
3. Corrales L, *et al.* 2012. *J. Immunol.* 189:4674. PubMed.

Description: CD88 is a single chain protein with seven membrane-spanning regions and a molecular weight of 43 kD. CD88 is expressed by neutrophils, eosinophils, and monocytes. There have been reports of CD88 expression in non-immune cells such as glial cells, cerebellar granule cells, cardiomyocytes, and vascular endothelial cells. CD88 is coupled to heterotrimeric G proteins such as G_i, and after C5a binding, the signal transduced by CD88 results in the activation of PLCβ, PI-3 kinase, and PLA₂, among other molecules, resulting in chemotaxis, granule enzyme release, and superoxide anion production.

Antigen References:

1. Schraufstatter IU, *et al.* 2009. *J. Immunol.* 182:3827.
2. Ippel JH, *et al.* 2009. *J. Biol. Chem.* 284:12363.
3. Monk PN, *et al.* 2007. *Br. J. Pharmacol.* 152:429.
4. Griffin RS, *et al.* 2007. *J. Neurosci.* 27:8699.
5. Lee H, *et al.* 2006. *Nat. Biotechnol.* 24:1279.
6. Buck E and Wells JA, *et al.* 2005. *P. Natl. Acad. Sci. USA* 102:2719.

Related Products:

Product
Purified Rat IgG2b, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
RTK4530

Application
FC, ICC, ICFC, IF, IHC, IP, WB
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.