

PerCP anti-mouse CD90.2 (Thy1.2)

Catalog # / Size: 140315 / 25 µg
140316 / 100 µg

Clone: 53-2.1

Isotype: Rat IgG2a, κ

Immunogen: Mouse thymus or spleen


Reactivity: Mouse, Does not react with Thy-1.1 (CD90.1)

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP under optimal conditions. The solution is free of unconjugated PerCP and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 mouse splenocytes stained with CD90.2 (53-2.1) PerCP

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.06 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP has a maximum absorption of 482 nm and a maximum emission of 675 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining¹ of frozen tissue section, immunofluorescence², and immunoprecipitation³.

Application References:

1. Aldrich M, *et al.* 2003. *J. Immunol.* 171:5562. (IHC)
2. Jameson J, *et al.* 2004. *J. Immunol.* 172:3573. (IF)
3. Okada C, *et al.* 1990. *J. Immunol.* 144:3473. (IP)

Description: CD90.2 is a 25-35 kD immunoglobulin superfamily member also known as Thy-1.2, a GPI-linked membrane molecule. It is expressed on hematopoietic stem cells and neurons, all thymocytes, and peripheral T cells in Thy1.2 bearing mouse strains (Balb/c, CBA/J, C3H/He, C57BL/-, DBA, NZB/-). CD90.2 is a glycosylphosphatidylinositol (GPI)-anchored membrane glycoprotein involved in signal transduction. CD90.2 is involved in costimulation of lymphocyte proliferation and induction of hematopoietic stem cells differentiation. CD90.2 has been shown to interact with CD45.

Antigen References:

1. Borrello M, *et al.* 1996. *Cell. Immunol.* 173:198.
2. Radrizzani M, *et al.* 1995. *J. Neurosci. Res.* 42:220.
3. Williams A, *et al.* 1982. *Science* 216:696.

Related Products:	Product	Clone	Application
	PerCP Rat IgG2a, κ Isotype Ctrl	RTK2758	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.