


Product Data Sheet

PerCP anti-rat CD90/mouse CD90.1 (Thy-1.1)

Catalog # / Size:	202512 / 100 µg
Clone:	OX-7
Isotype:	Mouse IgG1, κ
Immunogen:	Rat thymocyte Thy-1 antigen
Reactivity:	Rat, Mouse (AKR/J and PL mouse strains), Cross-Reactivity: Rabbit (Lapine), Guinea Pig
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PerCP under optimal conditions. The solution is free of unconjugated PerCP and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration:	0.2 mg/ml
Storage:	The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.



Lou rat thymocytes stained with OX-7

PerCP

Applications:

Applications: FC - Quality tested Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.06 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application. * PerCP has a maximum absorption of 482 nm and a maximum emission of 675 nm. Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemistry¹ of acetone-fixed frozen sections and zinc-fixed paraffin-embedded sections, immunoprecipitation², Western blotting², in vitro activation of leukocytes³, induction of endothelial cell permeability⁴, induction of glomerulonephritis⁵ in vivo. Application References: 1. Hermans MHA, et al. 1991. J. Histochem. Cytochem. 39:1627. (IHC) 2. Jeng CJ, et al. 1998. J. Cell Biol. 140:685. (IP, WB) 3. Nakashima I, et al. 1991. J. Immunol. 147:1153. 4. Ishizu A, et al. 1995. Int. Immunol. 7:1939. 5. Eitner F. 1997. Kidney. Int. 51:69. 6. Kawachi H, et al. 1992. Clin. Exp. Immunol. 88:399. 7. Dyer KD, et al. 2007. J. Immunol. 179:1693. (FC) PubMed 8. Hiramatsu Y, et al. 2010. J. Immunol. 87:703. (FC) PubMed Description: CD90, also known as Thy-1, is a 28-30 kD GPI-linked membrane glycoprotein. It is a member of the immunoglobulin superfamily and has been shown to interact with CD45 in signal transduction during lymphocyte proliferation and differentiation. CD90 is expressed on hematopoietic stem cells, neurons, thymocytes, peripheral T cells, fibroblasts, stromal cells. The OX-7 antibody reacts with rat CD90 and mouse CD90.1 (Thy-1.1) (which is expressed by mouse strains of AKR/J, PL, and FVB/N), but not mouse CD90.2. This antibody has been reported to induce leukocyte activation, vascular permeability, induce apoptosis in glomerular mesangial cells, and induce glomerulonephritis in vivo. Antigen References: 1. Campbell DG, et al. 1981. Biochem. J. 195:15. 2. Hosseinzadeh H, et al. 1993. J. Immunol. 150:1670. **Related Products: Product** Clone Application FC, ICFC FC, ICC, ICFC MOPC-21 PerCP Mouse IgG1, κ Isotype Ctrl Cell Staining Buffer RBC Lysis Buffer (10X)



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.