

LEAF™ Purified anti-mouse CD122 (IL-2R β)

Catalog # / Size: 123203 / 50 μ g
123204 / 500 μ g

Clone: TM- β 1

Isotype: Rat IgG2b, κ

Immunogen: Rat T cell line expressing mouse CD122 (IL-2R β)


Reactivity: Mouse

Preparation: The LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.

Formulation: 0.2 μ m filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.1 EU/ μ g of the protein (<0.01 ng/ μ g of the protein) as determined by the LAL test.

Concentration: 1.0 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. This LEAF™ solution contains no preservative; handle under aseptic conditions.



C57BL/6 mouse splenocytes stained with LEAF™ purified TM- β 1, followed by anti-rat IgG FITC

Applications:

Applications: FC - *Quality tested*
IP, Block, Depletion - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 μ g per 10^6 cells in 100 μ l volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation¹, blocking of IL-2 binding¹, and NK cell depletion² *in vivo*. The LEAF™ Purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for blocking of IL-2 binding *in vivo* and *in vitro* (Cat. No. 123204).

Application References:

1. Tanaka T, *et al.* 1991. *J. Immunol.* 147:2222.
2. Tanaka T, *et al.* 1993. *J. Exp. Med.* 178:1103.
3. Tanaka T, *et al.* 1992. *Int. Immunol.* 4:487.

Description: CD122 is a 70-75 kD IL-2 receptor β chain also known as IL-2R β , which is also shared by the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2R β chain can combine with either the common γ_c subunit (γ_c , CD132) alone or with the γ_c subunit and the IL-2R α subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The TM- β 1 antibody does inhibit IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human, CD8+ Tregs involved in the maintenance of T cell homeostasis.

Antigen References:

1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook Academic Press.*
2. Minami Y, *et al.* 1993. *Annu. Rev. Immunol.* 11:245.
3. Suzuki H, *et al.* 1995. *Science* 268:1472.
4. Shi Z, *et al.* 2009. *Eur. J. Immunol.* 39:2109.

Related Products: **Product**
LEAF™ Purified Rat IgG2b, κ Isotype Ctrl

Clone
RTK4530

Application
FC, ICFC, WB, IP, ICC, IF, IHC, FA



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.