

Product Data Sheet

APC anti-mouse CD117 (c-kit)

Catalog # / Size: 135107 / 25 µg
135108 / 100 µg

Clone: ACK2

Isotype: Rat IgG2b, κ

Immunogen: Murine IL-3 dependent mast cells


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 bone marrow cells stained with ACK2 APC and CD11b (M1/70) FITC


Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: ACK2 has been reported to block c-Kit function. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 135103). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 135114) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

Application References: 1. Yoshinaga K, *et al.* 1991. *Development* 113:689.
2. Broudy VC, *et al.* 1996. *Blood* 88:75.
3. Sachs N, *et al.* 2012. *PNAS*. 109:21468. PubMed.



C57BL/6 bone marrow cells stained with rat IgG2b APC isotype control and CD11b (M1/70) FITC

Description: CD117 is a 145 kD immunoglobulin superfamily member, also known as c-Kit and stem cell factor receptor (SCFR). It is a transmembrane tyrosine-kinase receptor that binds the c-Kit ligand (also known as steel factor, stem cell factor, and mast cell growth factor). CD117 is expressed on hematopoietic stem cells (including multipotent hematopoietic stem cells, progenitors committed to myeloid and/or erythroid lineages, and T and B cell precursors), mast cells, and acute myeloid leukemia (AML) cells. CD117 interaction with its ligand is critical for the development of hematopoietic stem cells.

Antigen References: 1. Barclay A *et al.* 1997. *The Leukocyte Antigen FactsBook Academic Press*.
2. Galli SJ. *et al.* 1994. *Adv. Immunol.* 55:1.
3. Ikuta K *et al.* 1992. *Annu. Rev. Immunol.* 10:759.
4. Besmer P *et al.* 1986. *Nature* 320:415.
5. Witte ON. *et al.* 1990. *Cell* 63:5.

Related Products:

Product
APC Rat IgG2b, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
TruStain fcX™ (anti-mouse CD16/32)

Clone
RTK4530

93

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.