

APC anti-mouse CD115 (CSF-1R)

Catalog # / Size: 135509 / 25 µg
135510 / 100 µg

Clone: AFS98

Isotype: Rat IgG2a, κ


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.


Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Thioglycolate-elicited Balb/c mouse peritoneal macrophages stained with CD11b FITC and AFS98 APC (top) or rat IgG2a,κ APC (bottom).



Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: Blocking of ligand binding¹. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays.

Application References:

1. Sudo T, *et al.* 1995. *Oncogene* 11:2469.
2. Murayama T, *et al.* 1999. *Circulation* 99:1740.
3. Jaeger BN, *et al.* 2012. *J. Exp. Med.* 209:565. PubMed

Description: CSF-1R, also known as CD115 and M-CSFR, is a single-pass type I membrane protein and member of the platelet-derived growth factor receptor family. This c-fms (Fms proto-oncogene) gene product's natural ligands include M-CSF and IL-34. Structural studies of CD115 have described an Ig-like extracellular domain, a transmembrane domain, an intracellular juxtamembrane domain, a split tyrosine kinase domain, and a C-terminal tail receptor. Receptor activation induces homodimerization in addition to phosphorylation and ubiquitination of intracellular residues. CD115 directly influences tissue macrophage and osteoclast differentiation and proliferation. It is expressed on monocytes/macrophages, peritoneal exudate cells, plasmacytoid and conventional dendritic cells, and osteoclasts.

Antigen References:

1. Sudo T, *et al.* 1995 *Oncogene* 11:2469.
2. Murayama T, *et al.* 1999 *Circulation* 99:1740.
3. Goswami S, *et al.* 2005 *Cancer Res.* 65:5278.
4. Yu W, *et al.* 2008 *J. Leuko. Biol.* 84:852.

Related Products:

Product	Clone	Application
APC Rat IgG2a, κ Isotype Ctrl	RTK2758	FC, ICFC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.