

Product Data Sheet

PE anti-mouse CD135

Catalog # / Size: 135305 / 50 µg
135306 / 200 µg

Clone: A2F10

Isotype: Rat IgG2a, κ

Immunogen: Mouse Flt3 transfected cell line


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**




Bone marrow cells from C57BL/6 mouse stained with A2F10 PE and CD45R/B220 (RA3-6B2) FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Sergejeva S, *et al.* 2004. *Blood* 103:1270.
2. Auffray C, *et al.* 2009. *J. Exp. Med.* 206:595.



Bone marrow cells from C57BL/6 mouse stained with rat IgG2a, κ PE and CD45R/B220 (RA3-6B2) FITC

Description: CD135, also known as Flk-2, Flt3, and Ly-72, is a type III tyrosine kinase receptor. It is expressed on early B lymphoid lineage cells in bone marrow, on primitive myeloid progenitors within the BM CD34+ cell population. Ligation of Flk-2 with Flt3 ligand regulates the growth of hematopoietic stem cells and promotes the survival of primitive hematopoietic progenitor cells with myeloid as well as B lymphoid potential. It was reported that the receptor tyrosine kinase Flt3 is required for dendritic cell development. Combined signaling through interleukin-7 receptors and Flt3 selectively promotes B-cell commitment and differentiation from uncommitted murine bone marrow progenitor cells.

Antigen References: 1. Waskow C, *et al.* *Nat. Immunol.* 9:676
2. Veiby OP, *et al.* 1996. *Blood* 88(4):1256
3. Veiby OP, *et al.* 1996. *J. Immunol.* 157(7):2953
4. Matthews W, *et al.* 1991. *Cell* 65(7):1143
5. Hannum C, *et al.* 1994. *Nature* 368(2):643
6. Ogawa M, *et al.* 1998. *Exp Hematol.* 26(6):478

Related Products:

Product
PE Rat IgG2a, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
TruStain fcX™ (anti-mouse CD16/32)

Clone
RTK2758

93

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.