

APC anti-mouse CD140b

Catalog # / Size: 136007 / 25 µg
136008 / 100 µg

Clone: APB5

Isotype: Rat IgG2a, κ

Immunogen: Mouse PDGFR-β-hlgG1 recombinant fusion protein


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Mouse fibroblast NIH/3T3 cells stained with APB5 APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported (for the relevant formats) applications include: Western blotting and blocking function². The LEAF™ purified antibody is recommended for functional assays.

Application References:

1. Sano H, *et al.* 2001. *Circulation* 103:2955.
2. Sano H, *et al.* 2002. *Am. J. Pathol.* 161:135. (Block)

Description: Platelet-derived growth factor receptor-β (PDGFR-β), CD140b, is one of two receptors for platelet-derived growth factors (PDGFs) and binds to all isoforms of PDGFs. PDGFRβ is a receptor tyrosine kinase that forms homodimers or heterodimers on the surface upon ligand binding and phosphorylates substrates. PDGFRs consist of either homodimers of α/α, β/β, or heterodimers of α/β. PDGF receptors, α and β, are single glycoproteins with intracellular tyrosine kinase domain. Their ligand, PDGF, is a mitogen for connective tissue and glial cells. CD140b is expressed on embryonic tissues and mesenchymal-derived cells adult mice. PDGF plays a role in wound healing and acts as a chemoattractant for fibroblasts, smooth muscle cells, glial cells, monocytes, and neutrophils.

Antigen References:

1. Soriano P, *et al.* 1994. *Genes Dev.* 8:1888
2. Takakura N, *et al.* 1996. *J Invest Dermatol.* 107:770
3. Yarden Y, *et al.* 1986. *Nature* 323:226

Related Products:Product

APC Rat IgG2a, κ Isotype Ctrl

Cell Staining Buffer

RBC Lysis Buffer (10X)

TruStain fcX™ (anti-mouse CD16/32)

Clone

RTK2758

93

Application

FC, ICFC

FC, ICC, ICFC

FC, ICFC

FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.