

Purified anti-mouse CD144 (VE-cadherin)

Catalog # / Size: 138001 / 50 µg
138002 / 500 µg

Clone: BV13

Isotype: Rat IgG1

Immunogen: VE-cadherin-Ig fusion protein


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Mouse endothelial cells b.End.3 stained with purified BV13 conjugated with Alexa Fluor® 647

Applications:

Applications: FC - *Quality tested*
WB, IF - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone BV13 recognizes an epitope between aa 45 and 56, and has a binding affinity of 5-15 nM.⁵ Additional reported applications (for relevant formats) include: Western blotting¹, blocking of cell interactions *in vivo*¹, and immunofluorescence microscopy⁴.

Application References:

1. Corada M, *et al.* 1999. *P. Natl. Acad. Sci. USA* 96:9815. (WB, Block)
2. Liao F, *et al.* 2000. *Cancer Res.* 60:6805. (FC)
3. Crosby CV, *et al.* 2005. *Blood* 105:2771. (FC)
4. Liao F, *et al.* 2002. *Cancer Res.* 62:2567. (IF)
5. May C, *et al.* 2005. *Blood* 105:4337. (epitope)

Description: CD144, also known as vascular endothelial-cadherin (VE-cadherin), is a 120 kD member of the type II Cadherin family. It is an endothelial specific hemophilic adhesion molecule involved in endothelial cell survival, migration, contact-dependent growth inhibition, and homophilic adhesion. VE-cadherin is essential for maintaining the integrity of the endothelial barrier *in vivo*.

Antigen References:

1. Allport JR, *et al.* 2002. *J. Leukocyte Biol.* 71:821.
2. Hirashima M, *et al.* 2009. *Blood* 93:1253.
3. Matsuyoshi N, *et al.* 1997. *Proc. Assoc. Am. Physicians* 109:362.
4. Matsumura K, *et al.* 2003. *Blood* 101:1367.
5. Hirashima M, *et al.* 2009. *Blood* 101:2261.
6. Gotsch U, *et al.* 1997. *J. Cell Sci.* 110:583.
7. Kataoka H, *et al.* 1997. *Dev. Growth Differ.* 39:729.

Related Products:

Product
Purified Rat IgG1, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
RTK2071

Application
FC, ICFC, ICC, IF, IHC, IP, WB
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.