


Anti-Human IgM PE

Catalog Number: 12-9998

For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD19 PerCP-Cy5.5 (cat. 45-0199) and Mouse IgG1 K Isotype Control PE (cat. 12-4714) (left) or Anti-Human IgM PE (cat. 12-9998) (right). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Human IgM PE

REF **Catalog Number:** 12-9998

Clone: SA-DA4

Concentration: 5 μ L (0.06 μ g)/test

Host/Isotype: Mouse IgG1, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Contains sodium azide



Description

The SA-DA4 monoclonal antibody reacts with the mu heavy chain of human IgM. It does not react with other classes of human immunoglobulin including IgD, IgG, or IgA. IgM is expressed intracellularly, during early stages of B lymphopoiesis, and then on the surface of mature B cells.

Our testing has revealed variations in staining of the non-B cell population that is predicted to be non-specific. This variation can be attributed to both donor variability and the anticoagulant used during blood draw. Therefore, we recommend using sodium citrate as the anticoagulant to minimize non-specific staining by this antibody.

Applications Reported

This SA-DA4 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This SA-DA4 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 μ L (0.06 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Szysko EA, Brun JG, Skarstein K, Peck AB, Jonsson R, Brokstad KA. Phenotypic diversity of peripheral blood plasma cells in primary Sjögren's syndrome. Scand J Immunol. 2011 Jan;73(1):18-28. (SA-DA4, FC)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Human IgM PE

Catalog Number: 12-9998

For Research Use Only. Not for use in diagnostic procedures.

Pack M, Trumpfheller C, Thomas D, Park CG, Granelli-Piperno A, Münz C, Steinman RM. DEC-205/CD205+ dendritic cells are abundant in the white pulp of the human spleen, including the border region between the red and white pulp. *Immunology*. 2008 Mar;123(3):438-46. (**SA-DA4**, IHC-fr)

Related Products

00-4222 Flow Cytometry Staining Buffer

12-4714 Mouse IgG1 K Isotype Control PE (P3.6.2.8.1)

45-0199 Anti-Human CD19 PerCP-Cy5.5 (HIB19)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com