


Anti-Human CD3 APC

Catalog Number: 17-0036

Also known as: Leu-4, T3

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD19 eFluor[®] 450 (cat. 48-0199) and Mouse IgG1 K Isotype Control APC (cat. 17-4714) (left) or Anti-Human CD3 APC (right). Cells in the lymphocyte gate were used for analysis.

Product Information



Contents: Anti-Human CD3 APC
Catalog Number: 17-0036
Clone: SK7
Concentration: 5 μ L (0.125 μ g)/test
Host/Isotype: Mouse IgG1, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer
Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.



Batch Code: Refer to vial



Use By: Refer to vial

Description

The SK7 monoclonal antibody reacts with human CD3e, a 20 kDa subunit of the TCR complex. Along with the other CD3 subunits γ and δ , the ϵ chain is required for proper assembly, trafficking and surface expression of the TCR complex. CD3 is expressed by thymocytes in a developmentally regulated manner and by all mature T cells. The SK7 and UCHT1 monoclonal antibodies cross-block binding, suggesting recognition of overlapping epitopes.

The antibody SK7 recognizes chimpanzee CD3.

Applications Reported

This SK7 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This SK7 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 μ L (0.125 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Kap YS, van Meurs M, van Driel N, Koopman G, Melief MJ, Brok HP, Laman JD, 't Hart BA. A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. *J Histochem Cytochem.* 2009 Dec;57(12):1159-67. (SK7.10, IHC frozen, chimpanzee, PubMed)

van Dongen JJ, Krissansen GW, Wolvers-Tettero IL, Comans-Bitter WM, Adriaansen HJ, Hooijkaas H, van Wering ER, Terhorst C. Cytoplasmic expression of the CD3 antigen as a diagnostic marker for immature T-cell malignancies. *Blood.* 1988 Mar;71(3):603-12

Related Products

17-4714 Mouse IgG1 K Isotype Control APC (P3.6.2.8.1)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Human CD3 APC

Catalog Number: 17-0036

Also known as: Leu-4, T3

RUO: For Research Use Only. Not for use in diagnostic procedures.

48-0199 Anti-Human CD19 eFluor[®] 450 (HIB19)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com