

## Purified anti-mouse CD146

**Catalog # / Size:** 134701 / 50 µg  
134702 / 200 µg

**Clone:** ME-9F1

**Isotype:** Rat IgG2a

**Immunogen:** Endothelial cell line TME-3H3

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.

## Applications:

**Applications:** FC - *Quality tested*

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:** 1. Schrage A, *et al.* 2008. *Histochem. Cell. Biol.* 129:441.

**Description:** CD146, also known as melanoma cell adhesion molecule (MCAM or Mel-CAM), MUC18, S-Endo1, and A32 antigen, is an integral membrane glycoprotein that belongs to the Ig superfamily. CD146 is strongly expressed by murine vascular endothelial cells. It is expressed on about 30% of neutrophils and 60% of NK cells. Unlike in humans, CD146 is undetectable on monocytes, dendritic cells, T cells, NKT cells, B cells, or smooth muscle cells in mouse. It has been reported that an increase in CD146 expression is associated with NK cell maturation. Combined with using CD27 and CD11b staining, CD146 may be an alternative marker to detect final stages of NK cell maturation and define NK cell subsets. CD146<sup>+</sup> NK cells were found to be less cytotoxic and to produce less IFN $\gamma$  than CD146<sup>-</sup> NK cells upon stimulation with target cells or activating antibodies. The role of CD146 on NK cell migration has yet to be investigated. The identification of CD146 ligand(s) will be crucial to address this issue.

**Antigen References:** 1. Despoix N, *et al.* 2008. *Eur. J. Immunol.* 38:2855.  
2. Sorrentino A, *et al.* 2008. *Exp. Hematol.* 36:1035.  
3. Bardin N, *et al.* 2009. *Arterioscler. Thromb. Vasc. Biol.* 29:746.

**Related Products:** **Product**


Purified Rat IgG2a,  $\kappa$  Isotype Ctrl  
RBC Lysis Buffer (10X)  
Cell Staining Buffer

**Clone**


RTK2758

**Application**

FC, ICC, IF, IHC, IP, WB  
FC, ICFC  
FC, ICC, ICFC



C57BL/6 splenocytes stained with ME-9F1 PE and NK-1.1 (PK136) APC



C57BL/6 splenocytes stained with rat IgG2a (RTK2758) PE isotype control and NK-1.1 (PK136) APC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.