


## Anti-Mouse CD8a APC

**Catalog Number:** 17-0081

**Also Known As:** CD8 alpha, Ly-2, Ly-35, Ly-B, Lyt-2

**RUO: For Research Use Only. Not for use in diagnostic procedures.**



Staining of mouse splenocytes with Anti-Mouse CD3 FITC (cat. 11-0031) and 0.06 ug of Rat IgG2a K Isotype Control APC (cat. 17-4321) (left) or 0.06 ug of Anti-Mouse CD8 APC (right). Total viable cells were used for analysis.

### Product Information

**Contents:** Anti-Mouse CD8a APC

**REF** **Catalog Number:** 17-0081

**Clone:** 53-6.7

**Concentration:** 0.2 mg/mL

**Host/Isotype:** Rat IgG2a, kappa

**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

**LOT** **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

### Description

The 53-6.7 monoclonal antibody reacts with the mouse CD8a molecule. CD8a is an approximately 32-34 kDa cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha beta) or as a homodimer (CD8 alpha alpha). A majority of thymocytes and a subpopulation of mature alpha beta TCR T cells express CD8 alpha beta while gamma delta TCR T cells, a subpopulation of intestinal intraepithelial lymphocytes (IELs) and dendritic cells express CD8 alpha alpha. CD8 binds to MHC class I and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells.

### Applications Reported

The 53-6.7 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

The 53-6.7 antibody has been tested by flow cytometric analysis of mouse thymocyte or splenocyte suspensions. This can be used at less than or equal to 0.125 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Mochimaru H, Usui T, Yaguchi T, Nagahama Y, Hasegawa G, Usui Y, Shimmura S, Tsubota K, Amano S, Kawakami Y, Ishida S. Suppression of alkali burn-induced corneal neovascularization by dendritic cell vaccination targeting VEGF receptor 2. *Invest Ophthalmol Vis Sci.* 2008 May;49(5):2172-7. (53-6.7, in vivo depletion, PubMed)

Yang Z, Day YJ, Toufektsian MC, Xu Y, Ramos SI, Marshall MA, French BA, Linden J. Myocardial infarct-sparing effect of adenosine A2A receptor activation is due to its action on CD4+ T lymphocytes. *Circulation.* 2006 Nov 7;114(19):2056-64. (53-6.7, in vivo depletion, PubMed)

Taylor JL, Ordway DJ, Trout J, Gonzalez-Juarrero M, Basaraba RJ, Orme IM. Factors associated with severe granulomatous pneumonia in Mycobacterium tuberculosis-infected mice vaccinated therapeutically with hsp65 DNA. *Infect Immun.* 2005 Aug;73(8):5189-93. (53-6.7, IHC frozen)

Grabbe S, Varga G, Beissert S, Steinert M, Pendl G, Seeliger S, Bloch W, Peters T, Schwarz T, Sunderkötter C, Scharffetter-Kochanek K. Beta2 integrins are required for skin homing of primed T cells but not for priming naïve T cells. *J Clin Invest.* 2002 Jan;109(2):183-92. (53-6.7, IHC frozen)

Ledbetter JA, Rouse RV, Micklem HS, Herzenberg LA. T cell subsets defined by expression of Lyt-1,2,3 and Thy-1 antigens. Two-parameter immunofluorescence and cytotoxicity analysis with monoclonal antibodies modifies current views. *J Exp Med.* 1980 Aug 1;152(2):280-95.

Ledbetter, J. A. and L. A. Herzenberg. Xenogeneic monoclonal antibodies to mouse lymphoid differentiation antigens. *Immunol Rev.* 1979;47:63-90.

**Related Products**

11-0031 Anti-Mouse CD3e FITC (145-2C11)

17-4321 Rat IgG2a K Isotype Control APC (eBR2a)

---

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)