


Anti-Mouse CD16/CD32 APC

Catalog Number: 17-0161

Also Known As:FCGR3, IGFR3; FCGR2, IGFR2; FC Receptor Block RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of BALB/c splenocytes with staining buffer (autofluorescence) (open histogram) or 0.125 ug of Anti-Mouse CD16/CD32 APC (filled histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD16/CD32 APC

REF Catalog Number: 17-0161

Clone: 93

Concentration: 0.2 mg/mL Host/Isotype: Rat IgG2a, lambda Formulation: aqueous buffer, 0.09% sodium azide, may

contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze.

Light sensitive material. LOT Batch Code: Refer to Vial Use By: Refer to Vial

Caution, contains Azide

Description

The 93 monoclonal antibody reacts with an epitope shared by mouse CD16 and CD32. CD16 (Fc gamma III Receptor) and CD32 (Fc gamma II Receptor) are the low affinity receptors for the mouse IgG Fc portion and are expressed by B cells, monocyte/macrophages, NK cells, and neutrophils.

Applications Reported

The 93 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The 93 antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Oliver, A. M., J. C. Grimaldi, et al. Independently ligating CD38 and Fc gammaRIIB relays a dominant negative signal to B cells.1999. Hybridoma 18(2): 113-9.

Related Products

17-4321 Rat IgG2a K Isotype Control APC (eBR2a)

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com