


Anti-Mouse CD19 APC

Catalog Number: 17-0193

RUO: For Research Use Only



Staining of BALB/c splenocytes with Anti-Mouse CD3e FITC (cat. 11-0031) and staining buffer (autofluorescence) (left) or 0.06 µg of Anti-Mouse CD19 APC (right). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Mouse CD19 APC


REF Catalog Number: 17-0193

Clone: eBio1D3 (1D3)


Concentration: 0.2 mg/ml


Host/Isotype: Rat IgG2a, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

LOT Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

The eBio1D3 monoclonal antibody reacts with mouse CD19, a 95 kDa transmembrane glycoprotein. CD19 is expressed by B cells during all stages of development excluding the terminally differentiated plasma cells. Follicular dendritic cells also express CD19. Together CD21, CD81, MHC class II, and CD19 form a multimolecular complex that associates with the BCR. Signaling through CD19 induces tyrosine phosphorylation, calcium flux and proliferation of B cells.

Applications Reported

This eBio1D3 (1D3) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBio1D3 (1D3) antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.125 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Cherukuri A, Cheng PC, Pierce SK. The role of the CD19/CD21 complex in B cell processing and presentation of complement-tagged antigens. *J Immunol.* 2001 Jul 1;167(1):163-72.

Inabe K, Kurosaki T. Tyrosine phosphorylation of B-cell adaptor for phosphoinositide 3-kinase is required for Akt activation in response to CD19 engagement. *Blood.* 2002 Jan 15;99(2):584-9.

Krop I, de Fougères AR, Hardy RR, Allison M, Schlissel MS, Fearon DT. Self-renewal of B-1 lymphocytes is dependent on CD19. *Eur J Immunol.* 1996 Jan;26(1):238-42. (1D3, FA, PubMed)

Shoham T, Rajapaksa R, Boucheix C, Rubinstein E, Poe JC, Tedder TF, Levy S. The tetraspanin CD81 regulates the expression of CD19 during B cell development in a postendoplasmic reticulum compartment. *J Immunol.* 2003 Oct 15;171(8):4062-72.

Krop I, Shaffer AL, Fearon DT, Schlissel MS. The signaling activity of murine CD19 is regulated during cell development. *J Immunol.* 1996 Jul 1;157(1):48-56. (1D3, FC, FA, PubMed)

Related Products

11-0031 Anti-Mouse CD3e FITC (145-2C11)

12-0112 Anti-Mouse CD11b PE (M1/70)
17-4321 Rat IgG2a K Isotype Control APC
47-0042 Anti-Mouse CD4 APC-eFluor® 780 (RM4-5)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com