


# **Anti-Human CD27 APC**

Catalog Number: 17-0279 Also Known As:TNFRSF7

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Mouse IgG1 K Isotype Control APC (cat. 17-4714) (open histogram) or Anti-Human CD27 APC (filled histogram). Cells in the lymphocyte gate were used for analysis.

### **Product Information**

Contents: Anti-Human CD27 APC
REF Catalog Number: 17-0279

Clone: O323

Concentration: 5 uL (0.25 ug)/test Host/Isotype: Mouse IgG1, kappa HLDA Workshop: IV T-186 Formulation: aqueous buffer, 0.09% sodium azide, may

contain carrier protein/stabilizer

**Temperature Limitation:** Store at 2-8°C. Do not freeze.

■ Light sensitive material.

■ Batch Code: Refer to Vial

■ Use By: Refer to Vial

↑ Caution, contains Azide

## Description

The O323 monoclonal antibody reacts with human CD27, a lymphocyte-specific member of the TNFR superfamily. CD27 is expressed by a subset of thymocytes and virtually all mature T cells and is upregulated upon T-cell stimulation. CD27 binds to CD70, and through this interaction, plays an important role in T cell-B cell interaction.

## **Applications Reported**

The O323 antibody has been reported for use in flow cytometric analysis.

### **Applications Tested**

This O323 antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 5  $\mu$ L (0.25  $\mu$ g) per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test.

### References

Knapp, W., B. Dorken, et al. eds. (1989). Leucocyte Typing IV: White Cell Differentiation Antigens. Oxford University Press. New York.

## **Related Products**

16-0271 Anti-CD27 Functional Grade Purified (LG.7F9) 17-4714 Mouse IgG1 K Isotype Control APC (P3.6.2.1)