


An Affymetrix Company


# **Anti-Human CD32 APC**

Catalog Number: 17-0329

Also known as: Low Affinity IgG Receptor 2a, FcGR2a, IGFR3

RUO: For Research Use Only. Not for use in diagnostic procedures.





Staining of normal human peripheral blood cells with Anti-Human CD19 eFluor® 450 (cat. 48-0199) and Mouse IgG1 K Isotype Control APC (cat. 17-4714) (left) or Anti-Human CD32 APC (right). Cells in the lymphocyte gate were used for analysis.

### **Product Information**

Contents: Anti-Human CD32 APC

REF Catalog Number: 17-0329 Clone: 6C4 (CD32)

Concentration: 5 uL (0.5 ug)/test Host/Isotype: Mouse IgG1, kappa



**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer **Temperature Limitation:** Store at 2-8°C. Do not

freeze. Light sensitive material. **Batch Code:** Refer to vial



Use By: Refer to vial Contains sodium azide



This 6C4 monoclonal antibody reacts with human CD32 (also known as Fc gammaRII). This antibody recognizes two isoforms of the receptor, Fc gammaRIIA and Fc gammaRIIB. These 40-kDa polymorphic transmembrane glycoproteins are expressed on B cells, granulocytes, monocytes, macrophages, and platelets. Moreover, these receptors are detected on natural killer cells. CD32 enables interaction between Fc gammaRII-expressing cells and opsonized antigen or IgG-containing immune complexes. Both receptors exhibit low affinity towards IgG and play a role in inflammation and autoimmune disease. This clone has been reported to inhibit Ig binding in a rosette blocking assay.

### **Applications Reported**

This 6C4 antibody has been reported for use in flow cytometric analysis.

### **Applications Tested**

This 6C4 (CD32) antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5  $\mu$ L (0.5  $\mu$ g) per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test.

#### References

Fossati G, Bucknall RC, and Edwards SW. Fcgamma receptors in autoimmune diseases. Eur. J. Clin. Invest. 2001 31(9):821-831.

Vely F, Gruel N, Moncuit J, Cochet O, Rouard H, Dare S, Galon J, Sautes C, Fridman WH, Teillaud JL. A new set of monoclonal antibodies against human Fc gamma RII (CD32) and Fc gamma RIII (CD16): characterization and use in various assays. Hybridoma 1997 Dec; 16(6):519-28. (6C4, FC, FA (blocking), Pubmed)

## **Related Products**

00-4222 Flow Cytometry Staining Buffer



An Affymetrix Company

# **Anti-Human CD32 APC**

Catalog Number: 17-0329

Also known as: Low Affinity IgG Receptor 2a, FcGR2a, IGFR3 RUO: For Research Use Only. Not for use in diagnostic procedures.

00-4300 10X RBC Lysis Buffer (Multi-species) 17-4714 Mouse IgG1 K Isotype Control APC (P3.6.2.8.1) 48-0199 Anti-Human CD19 eFluor® 450 (HIB19)