

Biotin anti-mouse CD200 (OX2)

Catalog # / Size: 123804 / 500 µg

Clone: OX-90

Isotype: Rat IgG2a, κ

Immunogen: Soluble fusion protein of the extracellular region of mouse OX-2 antigen with domains 3 and 4 of rat CD4 fusion protein.


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. **Do not freeze.**



C57BL/6 splenocytes stained with biotinylated OX-90, followed by Sav-PE

Applications:

Applications: FC - *Quality tested*
IHC - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: CD200 (OX-2 antigen) is a type-1 membrane glycoprotein containing two extracellular Ig-like domains. CD200 a highly conserved type I membrane glycoprotein that is expressed on a variety of cell types including thymocytes, some T cells, endothelial and follicular dendritic cells, B cells, and brain tissue (neurons); but not on NK cells, granulocytes, monocytes, or macrophages. CD200 costimulates T cell proliferation. It may regulate myeloid cell activity in a variety of tissues. CD200 is the ligand for CD200 receptor (CD200R). The CD200 Receptor is restricted to myeloid cells, and it is believed that its engagement with CD200 results in inhibition and/or downregulation of myeloid cell activity. Blocking of CD200/CD200R interactions decreases myeloid cell inhibitory thresholds which results in enhanced immune activation.

Antigen References:

1. Hoek RM, *et al.* 2000. *Science* 290:1768.
2. Gorczynski R, *et al.* 2004. *J. of Immunol.* 172:7744.
3. Gorczynski L, *et al.* 1999. *J. Immunol.* 162:774.
4. Rosenblum MD, *et al.* 2004. *Blood* 103:2691.
5. Zhang S, *et al.* 2004. *J. of Immunol.* 173:6786.
6. Barclay AN, *et al.* 2002. *Trends Immunol.* 23:285.

Related Products:

Product
Cell Staining Buffer
Biotin Rat IgG2a, κ Isotype Ctrl
TruStain fcX™ (anti-mouse CD16/32)

Clone

RTK2758
93

Application

FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.