

# Product Data Sheet

## Purified anti-mouse CD202b (Tie-2, CD202)

**Catalog # / Size:** 124001 / 50 µg  
124002 / 200 µg

**Clone:** TEK4

**Isotype:** Rat IgG1, κ


**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.



C57BL/6 bone marrow myeloid cells stained with purified TEK4, followed by biotinylated anti-rat IgG and Sav-PE

## Applications:

**Applications:** FC - *Quality tested*  
IF - *Reported in the literature*

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:** 1. Fathers KE, *et al.* 2005. *Am. J. Pathol.* 167:1753.  
2. Sanchez-Martin L, *et al.* 2011. *Blood* 117:88. PubMed

**Description:** CD202b, also known as Tie-2 or TEK, is a 145 kD type I transmembrane protein. It is a member of the receptor tyrosine kinase (RTK) family of proteins and is expressed by endothelial cells and their progenitors, quiescent hematopoietic stem cells (HSCs), and a subpopulation of monocytes. Angiopoietin-1 (Ang-1) is an activator of CD202b to promote, maintain, and stabilize mature vessels and to maintain HSCs in the quiescent state. Ang-2 is another ligand of CD202b, which is involved in postnatal angiogenesis and in antagonizing the effects of Ang-1. Tie-2 also binds to Ang-4.

**Antigen References:** 1. Fathers KE, *et al.* 2005. *Am. J. Pathol.* 167:1753.  
2. Yano M, *et al.* 1997. *Blood* 89:431726.  
3. Kolatsi-Joannou M, *et al.* 2001. *Dev. Dyn.* 222:1206.

**Related Products:** **Product**  
Cell Staining Buffer  
Purified Rat IgG1, κ Isotype Ctrl

**Clone**  
RTK2071

**Application**  
FC, ICC, ICFC  
FC, ICFC, ICC, IF, IHC, IP, WB



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biollegend.com/ordering#license](http://www.biollegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.