

PE anti-mouse CD223 (LAG-3)

Catalog # / Size: 125207 / 25 µg
125208 / 200 µg

Clone: C9B7W

Isotype: Rat IgG1, κ

Immunogen: Murine CD223-Ig fusion protein


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



ConA-stimulated C57BL/6 splenocytes stained with C9B7W PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

1. Workman CJ, *et al.* 2002. *Eur. Immunol.* 32:2255. (IP, ELISA)
2. Workman CJ, *et al.* 2005. *J. Immunol.* 174:688. (FC)
3. Li N, *et al.* 2004. *J. Immunol.* 173:6806. (IP, ELISA)

Description: CD223, also known as LAG-3, is 70 kD type I transmembrane glycoprotein within the Ig superfamily. It is the gene product of lymphocyte activation gene-3 (Lag3) with structural similarity to CD4. CD223 is expressed on activated T cells and NK cells, as well as T regulatory cells. CD223, like CD4, binds MHC class II molecules, but with higher affinity than CD4. In association with the CD3 complex, CD223 has been shown as a negative regulatory molecule to inhibit TCR-mediated signaling and regulate homeostatic T cell expansion. In contrast to inhibition, CD223 has an activation function to promote immune responses by inducing MHC class II signaling in antigen presenting cells.

Antigen References: 1. Baixeras E *et al.* 1992. *J. Exp. Med.* 176:327.

Related Products:

Product
Cell Staining Buffer
PE Rat IgG1, κ Isotype Ctrl
TruStain fcX™ (anti-mouse CD16/32)

Clone

RTK2071
93

Application

FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.