

FITC anti-mouse CD199 (CCR9)

Catalog # / Size: 128705 / 50 µg
128706 / 500 µg

Clone: CW-1.2

Isotype: Mouse IgG2a, κ


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 thymocytes stained with CW-1.2 FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: CCR9 is a member of the G protein coupled receptor family and is involved in T cell development in the thymus and in gut-associated immune response. Mouse CCR9 is highly expressed on different stages of thymocytes and upregulated on CD4+CD8+ cells. It is expressed to a much lower level on spleen and lymph node cells. Expression of CCR9 on γδ T cells in the intraepithelial and small intestine has been reported. The interaction of CCR9 with its ligand CCL25 (TECK, thymus-expressed chemokine) may direct the trafficking of developing T cells in the thymus and generation of gut-specific immunological memory.

Antigen References: 1. Zaballo A, *et al.* 1999. *J. Immunol.* 162:5671.
2. Wurbel MA, *et al.* 2007. *J. Immunol.* 178:7598.
3. Wurbel MA, *et al.* 2006. *Eur. J. Immunol.* 36:73.

Related Products:

Product
FITC Mouse IgG2a, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
TruStain fcX™ (anti-mouse CD16/32)

Clone
MOPC-173

93

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.