

Product Data Sheet

PE anti-mouse CD196 (CCR6)

Catalog # / Size: 129803 / 25 µg

129804 / 100 µg

Clone: 29-2L17

Isotype: Armenian Hamster IgG

Immunogen: N-terminal peptide (aa. 2-38) of mouse CCR6 fused with GST

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



C57BL/6 splenocytes stained RA3-6B2 (B220) FITC and 29-2L17

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For

immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is

recommended that the reagent be titrated for optimal performance for each application.

Description: CCR6 is a G-protein linked chemokine receptor which binds the chemokine CCL20/MIP-3a. It is expressed on B

lymphocytes and certain subsets of dendritic cells and T cells. CCR6 in reported to be involved in mucosal immune

93

response and lymphocyte migration and homeostasis.

1. Rossi D, *et al.* 1997. *J. Immunol.* 158:1033. 2. Lukacs N, *et al.* 2001. *J. Exp. Med.* 194:551. 3. Cook DN, *et al.* 2000. *Immunity* 12:495 **Antigen References:**

Application Related Products: Product Clone PE Armenian Hamster IgG Isotype Ctrl **HTK888**

Cell Staining Buffer RBC Lysis Buffer (10X)

TruStain fcX™ (anti-mouse CD16/32)

FC, ICFC FC, ICC, ICFC FC, ICFC



