

# Anti-Human/Mouse CD44 APC

Catalog Number: 17-0441 Also Known As:Pgp-1, MDU3, Hermes, Hyaluronate receptor RUO: For Research Use Only



Staining of C57BL/6 splenocytes with staining buffer (autofluorescence) (open histogram) or 0.03 µg of Anti-Human/Mouse CD44 APC (filled histogram). Total viable cells were used for analysis.

## Product Information

Contents: Anti-Human/Mouse CD44 APCFormulation: aqueous buffer, 0.09% sodium azide, may contain<br/>carrier protein/stabilizerREF Catalog Number: 17-0441<br/>Clone: IM7<br/>Concentration: 0.2 mg/ml<br/>Host/Isotype: Rat IgG2b, κFormulation: aqueous buffer, 0.09% sodium azide, may contain<br/>carrier protein/stabilizerImage: Concentration: 0.2 mg/ml<br/>Host/Isotype: Rat IgG2b, κFormulation: Store at 2-8°C. Do not freeze. Light<br/>sensitive material.Image: Concentration: 0.2 mg/ml<br/>Host/Isotype: Rat IgG2b, κImage: Concentration: Concentratic Concentration: Concent

### Description

The IM7 monoclonal antibody reacts with all isoforms of mouse CD44 (Pgp-1). CD44 is expressed by hematopoietic and non-hematopoietic cells. Bone marrow myeloid cells and memory T cells highly express this antigen and peripheral B and T cells can upregulate the expression of CD44. CD44 functions as an adhesion molecule through its binding to hyaluronate, an extracellular matrix component.

### Applications Reported

The IM7 antibody has been reported for use in flow cytometric analysis.

### **Applications Tested**

The IM7 antibody has been tested by flow cytometric analysis of mouse bone marrow cells and splenocyte suspensions. This can be used at less than or equal to 0.06  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Trowbridge, I. S., J. Lesley, et al. 1982.Biochemical characterization and cellular distribution of a polymorphic, murine cell-surface glycoprotein expressed on lymphoid tissues. Immunogenetics 15(3): 299-312.

Lesley, J. and I. S. Trowbridge 1982. Genetic characterization of a polymorphic murine cell-surface glycoprotein. Immunogenetics 15(3): 313-20. Maiti A, Maki G, Johnson P. TNF-alpha induction of CD44-mediated leukocyte adhesion by sulfation. Science. 1998. Oct 30;282(5390):941-3.

#### Related Products 17-4031 Rat IgG2b K Isotype Control APC

Not for further distribution without written consent. Copyright © 2000-2010 eBioscience, Inc. Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com