


Anti-Human CD69 APC

Catalog Number: 17-0699

Also Known As: Very Early Activation Antigen, VEA

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of 3-day PHA-stimulated normal human peripheral blood cells with Mouse IgG1 K Isotype Control APC (cat. 17-4714) (open histogram) or Anti-Human CD69 APC (filled histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Human CD69 APC REF Catalog Number: 17-0699

Clone: FN50

Concentration: 5 uL (0.015 ug)/test Host/Isotype: Mouse IgG1, kappa HLDA Workshop: IV A091 Formulation: aqueous buffer, 0.09% sodium azide, may

contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze.

■ Light sensitive material.

■ Batch Code: Refer to Vial

■ Use By: Refer to Vial

Caution, contains Azide

Description

The FN50 monoclonal antibody reacts with human CD69, also known as very early activation antigen (VEA). CD69 is approximately 30 kDa and is expressed on the cell-surface as a disulfide-linked dimer. CD69 is rapidly upregulated upon activation and expressed on lymphocytes, monocytes and platelets.

Applications Reported

This FN50 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This FN50 antibody has been pre-titrated and tested by flow cytometric analysis of resting and 6-hour TPA-activated human PBMC. This can be used at 5 μ L (0.015 μ g)/per test. A test is defined as the amount (μ g)/test of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

References

Schlossman, S., L. Bloumsell, et al. eds (1995). Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press. New York

Knapp, W., B. Dorken, et al. eds. (1989). Leucocyte Typing IV: White Cell Differentiation Antigens. Oxford University Press. New York.

Related Products

17-4714 Mouse IgG1 K Isotype Control APC (P3.6.2.1)