

Product Data Sheet

PE anti-mouse CD274 (B7-H1, PD-L1)

Catalog # / Size: 124307 / 25 µg
124308 / 100 µg

Clone: 10F.9G2

Isotype: Rat IgG2b, κ


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57/B6 mouse splenocytes were stained with anti-CD274 (clone 10F.9G2) PE (filled histogram) or rat IgG2b, κ PE (open histogram).

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence⁴ and blocking^{6,7,8,9}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 124303). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 124318) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

Application References:

1. Maier H, *et al.* 2007. *J. Immunol.* 178:2714.
2. Meng Q, *et al.* 2006. *Invest. Ophthalmol. Vis. Sci.* 47:4444. PubMed
3. Scarlett UK, *et al.* 2012. *J Exp Med.* 209:495. PubMed
4. Gracie N, *et al.* 2007. *Circulation* 116:2062. (IF)
5. Paterson AM, *et al.* 2011. *J. Immunol.* 187:1097.
6. Channappanavar R, *et al.* 2012. *PLoS One* 7:e39757. (Block)
7. Schreiber HA, *et al.* 2010. *PLoS One* 5:e11453. (Block) PubMed
8. Muthumani K, *et al.* 2011. *J. Immunol.* 187:2932. (Block) PubMed
9. Cripps JG, *et al.* 2010. *Hepatology* 52:1350. (Block) PubMed

Description: CD274, also known as B7-H1 or programmed death ligand 1 (PD-L1), is a 40 kD type I transmembrane protein and a member of the B7 family within the immunoglobulin receptor superfamily. It is expressed on T cells, B cells, NK cells, dendritic cells, IFN-γ activated endothelial cells, and monocytes. B7-H1 is one of the ligands of PD-1. The interaction of B7-H1 with PD-1 plays an important role in the inhibition of T cell responses. Other studies have shown that B7-H1 is able to costimulate T cell growth and cytokine production. CD274 is involved in costimulation essential for T lymphocyte proliferation and production of IL-10 and IFN-γ, in an IL-2-dependent and a PDCD1-independent manner. Its interaction with PDCD1 inhibits T-cell proliferation and cytokine production.

Antigen References:

1. Sharpe A, *et al.* 2007. *Nat. Immunol.* 8:239.
2. Dong H, *et al.* 1999. *Nat. Med.* 5:1365.
3. Freeman G, *et al.* 2000. *J. Exp. Med.* 192:1027.

Related Products:

Product
 Cell Staining Buffer
 PE Rat IgG2b, κ Isotype Ctrl
 TruStain fcX™ (anti-mouse CD16/32)

Clone

RTK4530
93

Application

FC, ICC, ICFC
 FC, ICFC
 FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.