

Product Data Sheet

Purified anti-mouse/human CD276 (B7-H3)

Catalog # / Size: 135602 / 100 µg

Clone: MIH35

Isotype: Rat IgG2a, κ

Immunogen: Mouse B7-H3 transfected L cell and P815


Reactivity: Mouse, **Cross-Reactivity:** Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Mouse B7-H3 transfected P815 cells stained with MIH35 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

1. Hashiguchi M, *et al.* 2008. *Proc Natl Acad Sci USA.* 105(30):10495.
2. del Rio ML, *et al.* 2011. *Transpl. Int.* 24:501. (FC) PubMed

Description: B7-H3 is a type I transmembrane protein belonging to the B7 family of co-stimulatory proteins. B7-H3 is mostly expressed on professional APCs including B cells, macrophages, and dendritic cells at low level. It is detected on various human and murine tumor cells, nasal and airway epithelial cells. Its expression on dendritic cells appears to be up-regulated by LPS. Initial studies have shown that B7-H3 provides a stimulatory signal to T cells. However, recent studies suggest a negative regulatory role for B7-H3 in T cell responses. Mouse B7-H3 protein inhibited T cell activation and effector cytokine production. Thus, the immunological function of B7-H3 remains unclear. B7-H3 is involved in the suppression of Th1-mediated immune responses and plays an important role in the development of pathogenic Th2 cells in a murine asthma model. Monoclonal antibody against B7-H3 enhances T cell proliferation in vitro and leads to exacerbated EAE in vivo. It was reported recently that the Triggering Receptor Expressed on Myeloid cells (TREM)-like Transcript 2 (TLT-2, TREML2) is a receptor for B7-H3 in mice. It remains controversial. Further studies are needed to identify the receptor of B7-H3.

Antigen References:

1. Nagashima O, *et al.* 2008. *J. Immunol.* 181:4062
2. Prasad DVR, *et al.* 2004. *J. Immunol.* 173:2500
3. Sun M, *et al.* 2002. *J. Immunol.* 168:6294
4. Xu J, *et al.* 2006. *Cellular and Molecular Immunology.* 3(3):235
5. Ford JW, *et al.* 2009. *Curr Opin Immunol.* 21(1):38
6. Leitner J, *et al.* 2009. *Eur. J. Immunol.* 2009. 39(7):1754

Related Products:

Product
Purified Rat IgG2a, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
RTK2758

Application
FC, ICC, IF, IHC, IP, WB
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.