

## **Product Data Sheet**

## Purified anti-mouse/human CD258 (LIGHT)

Catalog # / Size: 137102 / 500 µg

Clone: 15B2

**Isotype:** Mouse IgG1,  $\kappa$ 

Immunogen: Recombinant extracellular domain of murine LIGHT

Reactivity: Mouse, Human<sup>1</sup>

**Preparation:** The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.

## **Applications:**

**Applications:** ELISA - Quality tested IP - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by ELISA. For ELISA, the suggested use of this reagent is 10 µg/ml.

It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for relevant formats) include: in vivo blocking of LIGHT activity in a DSS-induced

acute intestinal inflammation model<sup>1</sup>; in vitro blocking of the LIGHT-induced expression of CXCL8/CXCL2 by BFS-1

and A375 cells<sup>1</sup>, and immunoprecipitation<sup>1</sup>.

Application References: 1. Jungbeck M, et al. 2009. J. Immunol. 128:451.

Description: CD258, also known as LIGHT, belongs to the tumor necrosis factor (TNF) superfamily. CD258 is a homotrimer

composed of type II single-pass chains. The cleavage of this molecule from the cell surface results in a soluble form of LIGHT. CD258 stimulates T cell proliferation, CTL activity, Th1 cytokine production and induces apoptosis in some tumor lines. CD258 is expressed by T cells, NK cells, macrophages and platelets. The receptors of LIGHT are HVEM,

LTβR and DCR3. CD258 has an important role in the cell-mediated immune response and in graft versus host

disease.

**Antigen References:** 1. Tama K, et al. 2000. Nat. Med. 6:283. 2. Cohavey O, et al. 2005. J. Immunol. 174:646.

3. Madge LA, et al. 2008. J. Immunol. 180:3467.

**Related Products: Product** Clone Application

RBC Lysis Buffer (10X)

FC, ICFC FC, ICC, ICFC Cell Staining Buffer

Purified Mouse IgG1, κ Isotype Ctrl MOPC-21 FC, ICFC, ICC, IF, IHC, IP, WB



