

PE anti-mouse CD254 (TRANCE, RANKL)

Catalog # / Size: 510005 / 50 µg
510006 / 200 µg

Clone: IK22/5

Isotype: Rat IgG2a, κ

Immunogen: NSO-derived recombinant mouse TRANCE


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



TRANCE transfected cells stained with IK22-5 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric analysis, the suggested use of this reagent is ≤ 0.25 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: The IK22/5 antibody has been reported to block the binding of RANK ligand-Fc to RANK. Additional reported applications (for the relevant formats) include: immunoprecipitation², Western blotting², and blocking of ligand binding^{1,2,4}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 510008).

Application References:

1. Miyahira Y, *et al.* 2003. *J. Immunol.* 171:6344. (Block)
2. Personal communication (Block IP WB)
3. Gao Y, *et al.* 2007. *J. Clin. Invest.* 117:122.
4. Kamijo S, *et al.* 2006. *Biochem. Biophys. Res. Commun.* 347:124. (Block)
5. Fionda C, *et al.* 2007. *J. Immunol.* 178:4039.
6. Haslam SZ, *et al.* 2008. *Endocrinology.* 149:2098. (Block) PubMed

Description: CD254 is a 19 kD TNF superfamily member also known as TRANCE (TNF-related activation induced cytokine), RANK ligand, RANKL, TNFSF11, OPGL, and ODF. TRANCE is expressed on activated T cells and osteoclasts and has been implicated in the regulation of T cell and dendritic cell interactions as well as osteoclast differentiation.

Antigen References:

1. Fitzgerald K, *et al.* Eds. 2001. *The Cytokine FactsBook.* Academic Press San Diego.
2. Josien R, *et al.* 1999. *J. Immunol.* 162:2562.
3. Takayanagi H, *et al.* 2000. *Nature* 408:600.
4. Wong B, *et al.* 1999. *Mol. Cell.* 4:1041.

Related Products:	Product	Clone	Application
	Cell Staining Buffer		FC, ICC, ICFC
	Fixation Buffer		ICC, ICFC
	Permeabilization Wash Buffer (10X)		ICC, ICFC, IHC
	Brefeldin A Solution (1,000X)		ICFC
	Monensin Solution (1,000X)		ICFC
	PE Rat IgG2a, κ Isotype Ctrl	RTK2758	FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.