


Anti-Human CD161 APC

Catalog Number: 17-1619

Also known as: NKR-P1A, NKR-P1A

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD4 eFluor® 450 (cat. 48-0048) and Mouse IgG1 K Isotype Control APC (cat. 17-4714) (left) or Anti-Human CD161 APC (right). Cells in the lymphocyte gate were used for analysis.

Product Information



Contents: Anti-Human CD161 APC

Catalog Number: 17-1619

Clone: HP-3G10

Concentration: 5 µL (0.25 µg)/test

Host/Isotype: Mouse IgG1



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Description

This HP-3G10 monoclonal antibody reacts with CD161 (also known as NKR-P1A), a member of the C-type lectin superfamily. The human homologue of NK1.1 in C57BL/6 mice, CD161 is expressed on natural killer cells and is upregulated in response to IL-12. CD161 can also be detected on various T cell subsets, including memory/effector CD4+ T cells, CD8+ T cells, gamma delta TCR T cells, and a subset of CD3+ thymocytes. Finally, CD161 expression has been demonstrated on human Th17 CD4+ T cells. The function of this receptor is unclear although studies suggest a possible stimulatory role. Nevertheless, Lectin-like transcript-1 (LLT1), which is also known as osteoclast inhibitory lectin, has been identified as the ligand for CD161.

Applications Reported

This HP-3G10 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This HP-3G10 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 µL (0.25 µg) per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

References

Cossmi, L, et al. Human interleukin 17-producing cells originate from a CD161+CD4+ T cell precursor. J. Exp. Med. 2008; 205(8): 1903-16.

Rosen DB, Cao W, Avery DT, Tangye SG, Liu YJ, Houchins JP, Lanier LL. Functional consequences of interactions between human NKR-P1A and its ligand LLT1 expressed on activated dendritic cells and B cells. J. Immunol. 2008 May 15; 180(10):6508-17.

Márquez C, Trigueros C, Franco JM, Ramiro AR, Carrasco YR, López-Botet M, Toribio ML. Identification of a common developmental pathway for thymic natural killer cells and dendritic cells. Blood. 1998 Apr 15;91(8):2760-71. (HP-

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Human CD161 APC

Catalog Number: 17-1619

Also known as: NKR-P1A, NKRP1A

RUO: For Research Use Only. Not for use in diagnostic procedures.

3G10, FC, Pubmed)

Related Products

17-4714 Mouse IgG1 K Isotype Control APC (P3.6.2.8.1)

48-0048 Anti-Human CD4 eFluor® 450 (OKT4 (OKT-4))