


Anti-Mouse CD317 (BST2, PDCA-1) APC

Catalog Number: 17-3172

Also known as: plasmacytoid DC Ag-1

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of SJL splenocytes with Anti-Human/Mouse CD45R (B220) FITC (cat. 11-0452) and 0.125 ug of Rat IgG2b K Isotype Control APC (cat. 17-4031) (left) or 0.125 ug of Anti-Mouse CD317 (BST2, PDCA-1) APC (right). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD317 (BST2, PDCA-1) APC



Catalog Number: 17-3172

Clone: eBio927

Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2b, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. DO NOT FREEZE. LIGHT-SENSITIVE MATERIAL.

Batch Code: Refer to vial

Use By: Refer to vial



LOT



Description

The eBio927 monoclonal antibody reacts with PDCA-1 (BST2, CD317), a specific marker of plasmacytoid dendritic cells (pDC), also known as type I IFN-producing cells (IPC) in the naïve mouse. Mouse IPCs are typically CD11c+, CD11b-, B220+, Ly-6C+, and CD62L+. PDCA-1 is predominantly expressed by IPCs in the naïve mouse which represents a very minor population (<0.5%) of splenocytes. It is upregulated on numerous cell types following stimulation which triggers an IFN response. PDCA-1 cycles between cell surface and intracellular compartments and may function to regulate trafficking of secreted cytokines. PDCA-1 (BST2) is the protein recognized by antibody 120G8.

The eBio927 monoclonal antibody has also been shown to have functional activity. The epitope recognized by eBio927 is distinct from eBio129c; thus, the antibodies can be used to costain, purify and identify pDCs.

Applications Reported

This eBio927 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBio927 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Blasius A.L., E. Giurisat, M. Cella, R.D. Schreiber, A.S. Shaw, M. Colonna. 2006. Bone Marrow Stromal Cell Antigen 2 Is a Specific Marker of Type I IFN-Producing Cells in the Naïve Mouse, but a Promiscuous Cell Surface Antigen Following IFN Stimulation. J Immunol. 177:3260-3265 (927, FC, FA)

Related Products

11-0452 Anti-Human/Mouse CD45R (B220) FITC (RA3-6B2)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Mouse CD317 (BST2, PDCA-1) APC

Catalog Number: 17-3172

Also known as: plasmacytoid DC Ag-1

RUO: For Research Use Only. Not for use in diagnostic procedures.

17-4031 Rat IgG2b K Isotype Control APC

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com