

APC anti-mouse CD309 (VEGFR2, Flk-1)

Catalog # / Size: 136405 / 25 µg
136406 / 100 µg

Clone: Avas12

Isotype: Rat IgG2a, κ

Immunogen: Murine Flk1 fused to hlgG Fc


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



bEnd.3 endothelial cells stained with Avas12a1 APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Avas12 recognizes a different epitope than clone 89B3A5.

Application References: 1. Kataoka H, *et al.* 1997. *Dev. Growth Differ.* 39:729.

Description: CD309 is also known as vascular endothelial growth factor receptor 2 (VEGFR2) and fetal liver kinase-1 (Flk-1). CD309 is a member of the tyrosine protein kinase family that contains a single pass transmembrane receptor with a protein kinase domain and seven immunoglobulin-like domains in the extracellular region. CD309 is expressed at high levels in adult heart, lung, kidney, brain, and skeletal muscle. It's a receptor for VEGF or VEGFC, and plays an important role in the development of vascular endothelial cells, hematopoietic cells, and vascular permeability.

Antigen References: 1. Kaburn N, *et al.* 1997. *Development.* 124:2039
2. Patterson C, *et al.* 1995. *J. Bio. Chem.* 270:23111
3. Nishikawa SI, *et al.* 1998. *Immunity* 8 (6):761
4. Shalaby F, *et al.* 1997. *Cell.* 89 (6):981

Related Products:	Product	Clone	Application
	APC Rat IgG2a, κ Isotype Ctrl	RTK2758	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.