


# **Anti-Mouse IgM APC**

Catalog Number: 17-5790

Also known as: Immunoglobulin M

RUO: For Research Use Only. Not for use in diagnostic procedures.



## Description

The II/41 monoclonal antibody reacts with the  $\mu$  heavy chain of mouse IgM. It does not react with other classes of mouse immunoglobulin including IgD, IgG or IgA. IgM is expressed intracellularly, during early stages of B lymphopoiesis, and then on the surface of more mature B cells in the bone marrow and peripheral B cells. Fluorochrome conjugated II/41 can be used as a detection secondary for mouse IgM.

#### **Applications Reported**

The II/41 antibody has been reported for use in flow cytometric analysis.

#### **Applications Tested**

The II/41 antibody has been tested by flow cytometric analysis of mouse bone marrow cells. This can be used at less than or equal to 0.25  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

#### References

Laszlo G, Hathcock KS, Dickler HB, Hodes RJ. 1993. Characterization of a novel cell-surface molecule expressed on subpopulations of activated T and B cells. J Immunol. 150(12):5252-62.

## **Related Products**

11-0452 Anti-Human/Mouse CD45R (B220) FITC (RA3-6B2) 17-4321 Rat IgG2a K Isotype Control APC (eBR2a)