


Anti-Mouse Ly-6A/E (Sca-1) APC

Catalog Number: 17-5981

Also Known As: Sca1, Ly6AE, Ly6A, Ly6E

RUO: For Research Use Only



Staining of unstimulated (left) and 4-day ConA-stimulated (right) BALB/c splenocytes with 0.03 µg of Rat IgG2a κ Isotype Control APC (cat. 17-4321) (open histogram) or 0.03 µg of Anti-Mouse Ly-6A/E (Sca-1) APC (filled histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse Ly-6A/E (Sca-1) APC

 Catalog Number: 17-5981

Clone: D7

Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2a, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The D7 monoclonal antibody reacts with mouse Sca-1, an 18 kDa member of the Ly-6 family of GPI-linked surface proteins. D7 reacts with both Ly-6E.1 and Ly-6A.2 molecules expressed by mouse hematopoietic stem cells, myeloid population and peripheral T and B cells. Non-activated splenocytes from Ly-6.2 strains including C57BL, SJL, 129, AKR, and others demonstrate higher frequency of D7-positive cells compared to the Ly-6.1 strains such as BALB/c, C3H, NZB and others. Sca-1 is expressed at high levels upon activation regardless of the Ly-6 haplotype. The Ly-6 family is involved in regulation and function of T cell activation. Sca-1 is a major phenotypic marker for mouse hematopoietic progenitor/stem cell subset.

Applications Reported

D7 has been reported for use in flow cytometric analysis.

Applications Tested

The D7 antibody has been tested by flow cytometric analysis of mouse splenocyte and bone marrow cell suspensions and can be used at less than or equal to 0.06 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Ortega, G., P. E. Kory, et al. (1986). "Role of Ly-6 in lymphocyte activation. I. Characterization of a monoclonal antibody to a nonpolymorphic Ly-6 specificity." *J Immunol* 137(10): 3240-6.

Malek, T. R., G. Ortega, et al. (1986). "Role of Ly-6 in lymphocyte activation. II. Induction of T cell activation by monoclonal anti-Ly-6 antibodies." *J Exp Med* 164(3): 709-22.

Related Products

17-4321 Rat IgG2a K Isotype Control APC

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com