

Product Data Sheet

LEAF™ Purified anti-mouse F4/80

Catalog # / Size: 123103 / 50 µg
123104 / 500 µg

Clone: BM8

Isotype: Rat IgG2a, κ

Immunogen: Murine macrophages


Reactivity: Mouse

Preparation: The LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.

Formulation: 0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.1 EU/µg of the protein (<0.01 ng/µg of the protein) as determined by the LAL test.

Concentration: 1.0 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. This LEAF™ solution contains no preservative; handle under aseptic conditions.



Thioglycollate elicited Balb/c peritoneal macrophage cells stained with LEAF™ purified BM8, followed by anti-rat IgG FITC

Applications:

Applications: FC - *Quality tested*
IHC, WB - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections^{1,2} and Western blotting.

Application References:

- Schaller E, *et al.* 2002. *Mol. Cell. Biol.* 22:8035. (IHC)
- Stevceva L, *et al.* 2001. *BMC Clin Pathol.* 1:3. (IHC)
- Kobayashi M, *et al.* 2008. *J. Leukocyte Biol.* 83:1354. PubMed
- Poeckel D, *et al.* 2009. *J. Biol Chem.* 284:21077. PubMed

Description: F4/80 is a 160 kD glycoprotein. It is characterized as a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has been widely used as a murine macrophage marker, which is expressed on majority of tissue macrophages including peritoneal macrophages, macrophages in lung, gut, thymus and red pulp of spleen (but not on the macrophages located in T cell areas of the spleen, lymph node and Peyer's patch), Kuffer cells, Langerhans cells, bone marrow stromal cells. F4/80 has also been shown on a subset of dendritic cells. The biological ligand of F4/80 has not been identified, but it has been reported that F4/80 is required for induction of CD8 T cells-mediated peripheral tolerance.

Antigen References:

- Austy JM and Gordon S. 1981. *Eur. J. Immunol.* 11:805.
- Hume DA, *et al.* 1983. *J. Exp. Med.* 158:1522.
- Ruedl C, *et al.* 1996. *Eur. J. Immunol.* 26:1801.
- McKnight AJ, *et al.* 1996. *J. Biol. Chem.* 271:486.
- Lin HH, *et al.* 2005. *J. Exp. Med.* 201:1615.

Related Products:

Product
APC Goat anti-rat IgG (minimal x-reactivity)
FITC Goat anti-rat IgG (minimal x-reactivity)
HRP Goat anti-rat IgG (minimal x-reactivity)
PE Goat anti-rat IgG (minimal x-reactivity)
Cell Staining Buffer
LEAF™ Purified Rat IgG2a, κ Isotype Ctrl

Clone
Poly4054
Poly4054
Poly4054
Poly4054
RTK2758

Application

FC
FC
ELISA, IF, IHC, WB
FC
FC, ICC, ICFC
FC, ICFC, WB, IP, ICC, IF, IHC, FA



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.