

PE anti-mouse DC Marker (33D1)

Catalog # / Size: 124905 / 50 µg

Clone: 33D1

Isotype: Rat IgG2b, κ

Immunogen: Mouse dendritic cells


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 splenocytes were stained with I-A/I-E APC and DC Marker (clone 33D1) PE (top) or rat IgG2b, κ PE isotype control (bottom).

Applications:


Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Crowley MT, et al. 1990. *J. Immunol. Methods* 133:55.
2. Nussenzweig MC, et al. 1982. *P. Natl. Acad. Sci. USA* 79:161.

Description: 33D1 antigen, also known as DCIR2 (dendritic cell inhibitory receptor 2), is reported to be a dendritic cell marker on subpopulation DC in spleen, thymus, Peyer's patch. It is reported that most N418 (CD11c)+ cells in spleen are also 33D1 positive. Bone marrow dendritic cells upregulate 33D1 antigen expression after incubation with GM-CSF.

Related Products:	Product	Clone	Application
	PE Rat IgG2b, κ Isotype Ctrl	RTK4530	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.