

## **Product Data Sheet**

## **FITC anti-mouse GITR**

Catalog # / Size: 126307 / 50 µg

126308 / 200 µg

Clone: DTA-1

**Isotype:** Rat IgG2b,  $\lambda$ 

Immunogen: mouse CD25+CD4+ T cells

Reactivity: mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with


FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For

immunofluorescent staining, the suggested use of this reagent is ≤1.0 μg per million cells in 100 μl volume. It is

recommended that the reagent be titrated for optimal performance for each application.

Description: GITR glucocorticoid-induced TNFR-related gene, is a member of the TNF receptor superfamily, also known as

TNFRŠF18,and AITR (in humans).It is expressed at low levels on resting T lymphocytes and at high levels on CD25+CD4+Treg cells. The expression of GITR on T cells can be upregulated upon activation. Interaction of GITR with its ligand (GITRL) has been demonstrated to augment T cell activation, proliferation, cytokine production, as well as MAPKs and NF-kB activation, and abrogate the inhibitory function of CD25+CD4+ T reg cells.In vivo activation of

GITR causes development of autoimmune diseases and restores the suppressed immune response.

1. Tone M, et al. 2003. Proc. Natl. Acad. Sci. USA 100:15059 Antigen References:

2. Shimzu J et al. 2002 Nat Immunol 3:135

3. Stephens GL *et al.* 2004. *J. Immunol.* 173:5008 4. McHugh RS, *et al.* 2002. *Immunity* 16:311

**Related Products: Product** Clone Application TruStain fcX™ (anti-mouse CD16/32)







