

Purified anti-mouse B7-H4 (B7S1, B7X)

Catalog # / Size: 139402 / 100 µg

Clone: HMH4-5G1

Isotype: Armenian Hamster IgG

Immunogen: Mouse B7-H4 human IgG1 Fc fusion protein


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Mouse B7-H4 transfected 293 cells stained with purified HMH4-5G1 conjugated with PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.06 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported application (for relevant formats) include: blocking¹.

Application References: 1. Personal communication (Block)

Description: B7-H4 is a newly discovered B7 family member that negatively regulates T cell immunity. *In vitro*, B7-H4 inhibits CD4+ and CD8+ T cell proliferation, cytokine production, and generation of alloreactive cytotoxic T-cells (CTLs). *In vivo*, blockage of endogenous B7-H4 by specific monoclonal antibodies promotes T cell responses. A new reported function of B7-H4 is as an important negative regulator of innate immunity through growth inhibition of neutrophils. It has been reported that B7-H4 is not constitutively expressed on peripheral tissues but can be induced to express on T cells, B cells, macrophages, and dendritic cells. B7-H4 is expressed on some tumor cancer cells. The role of B7-H4 in tumor progression may be to transform precancerous cells and then protect them from immunosurveillance. Although B- and T-lymphocyte attenuator (BTLA) was proposed to be the receptor for B7-H4, further studies are needed to identify the inhibitory receptor of B7-H4.

Antigen References: 1. Choi IH, *et al.* 2003. *J. Immunol.* 171:4650.
2. Prasad DV, *et al.* 2003. *Immunity* 18:863.
3. Kryczek I, *et al.* 2006. *J. Exp. Med.* 203:871.

Related Products:

Product
Purified Armenian Hamster IgG Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
HTK888

Application
FC, ICC, ICFC, IF, IP, WB
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.