


Anti-Mouse TNF alpha APC

Catalog Number: 17-7321

Also Known As: Tumor Necrosis Factor alpha

RUO: For Research Use Only



Intracellular staining of Mouse Cytokine Positive Control Cells (cat. 00-4500) with Anti-Mouse CD4 PE (cat. 12-0041) and 0.125 ug of Rat IgG1 K Isotype Control APC (cat. 17-4301) (left) or 0.125 ug of Anti-Mouse TNF alpha APC (right).

Product Information

Contents: Anti-Mouse TNF alpha APC


REF **Catalog Number:** 17-7321

Clone: MP6-XT22

Concentration: 0.2 mg/mL


Host/Isotype: Rat IgG1, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

LOT **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

Description

The MP6-XT22 antibody reacts with mouse tumor necrosis factor-alpha (TNF alpha), a 17 kDa cytokine produced by monocytes, macrophages, neutrophils, NK cells and CD4(+) T cells. TNF alpha has cytolytic activity against a range of tumor cells and is important in immune regulation. TNF alpha forms dimers and trimers and also exists as a 26 kDa membrane-bound form.

Applications Reported

The MP6-XT22 antibody has been reported for use intracellular staining for flow cytometric analysis.

Applications Tested

The MP6-XT22 antibody has been tested by intracellular staining with flow cytometric analysis.. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Chackerian B, Lowy DR and Schiller JT. Conjugation of a self-antigen to papillomavirus-like particles allows for efficient induction of protective autoantibodies. *J Clin Invest.* 2001. 108(3):415-23. (IHC frozen, PubMed)

Williams RO, Mauri C, et al. Therapeutic actions of cyclosporine and anti-tumor necrosis factor alpha in collagen-induced arthritis and the effect of combination therapy. *Arthritis Rheum.* 1998. 41(10):1806-12. (IHC frozen, PubMed)

Hunter CA, Litton MJ, et al. Immunocytochemical detection of cytokines in the lymph nodes and brains of mice resistant or susceptible to toxoplasmic encephalitis. *J Infect Dis.* 1994. 170(4): 939-45.

Litton MJ, Sander B, et al. Early expression of cytokines in lymph nodes after treatment in vivo with Staphylococcus enterotoxin B. *J Immunol Methods* 1994. 175(1): 47-58.

Abrams JS, Roncarolo MG, et al. Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. *Immunol Rev.* 1992. 127: 5-24.

Related Products

00-4500 Mouse Cytokine Positive Control Cells

00-5523 Foxp3 Staining Buffer Set
12-0041 Anti-Mouse CD4 PE (GK1.5)
17-4301 Rat IgG1 K Isotype Control APC
88-8823 Fixation & Permeabilization Buffers

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com