


## Anti-Mouse CD279 (PD-1) APC

**Catalog Number:** 17-9981

**Also known as:** Pdc1, Programmed cell death protein 1

**RUO: For Research Use Only. Not for use in diagnostic procedures.**



Staining of unstimulated (orange) or ConA-stimulated (purple) splenocytes with 0.5  $\mu$ g of Anti-Mouse CD279 (PD-1) APC. Total viable cells were used for analysis.

### Product Information



**Contents:** Anti-Mouse CD279 (PD-1) APC

**Catalog Number:** 17-9981

**Clone:** RMP1-30

**Concentration:** 0.2 mg/mL

**Host/Isotype:** Rat IgG2b, kappa



**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

**Temperature Limitation:** Store at 2-8°C. Do not freeze. Light-sensitive material.

**Batch Code:** Refer to vial

**Use By:** Refer to vial

**Contains sodium azide**



### Description

The RMP1-30 antibody reacts with mouse PD-1 (programmed death-1), a 55 kDa member of the Ig superfamily. PD-1 contains the immunoreceptor tyrosine-based inhibitory motif (ITIM) and plays a key role in peripheral tolerance and autoimmune disease in mice. PD-1 is expressed mainly on activated T and B lymphocytes. Two novel B7 Family members have been identified as PD-1 ligands, PD-L1 (B7-H1) and PD-L2 (B7-DC). Evidence reported to date suggests overlapping functions for these ligands and their constitutive expression on some normal tissues and upregulation on activated antigen-presenting cells. RMP1-30 does not block the binding of either B7-H1-Ig or B7-DC-Ig to PD-1 transfectants.

### Applications Reported

This RMP1-30 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

This RMP1-30 antibody has been tested by flow cytometric analysis of stimulated mouse splenocytes. This can be used at less than or equal to 1  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from  $10^5$  to  $10^8$  cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Matsumoto K, Inoue H, Nakano T, Tsuda M, Yoshiura Y, Fukuyama S, Tsushima F, Hoshino T, Aizawa H, Akiba H, Pardoll D, Hara N, Yagita H, Azuma M, Nakanishi Y. B7-DC regulates asthmatic response by an IFN-gamma-dependent mechanism. *J Immunol.* 2004 Feb 15;172(4):2530-41.

Nishimura H, Okazaki T, Tanaka Y, Nakatani K, Hara M, Matsumori A, Sasayama S, Mizoguchi A, Hiai H, Minato N, Honjo T. Autoimmune dilated cardiomyopathy in PD-1 receptor-deficient mice. *Science.* 2001 Jan 12;291(5502):319-22.

Freeman GJ, Long AJ, Iwai Y, Bourque K, Chernova T, Nishimura H, Fitz LJ, Malenkovich N, Okazaki T, Byrne MC,

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.ebioscience.com](http://www.ebioscience.com) •  
[info@ebioscience.com](mailto:info@ebioscience.com)

---

## Anti-Mouse CD279 (PD-1) APC

**Catalog Number:** 17-9981

**Also known as:** Pcd1, Programmed cell death protein 1

**RUO: For Research Use Only. Not for use in diagnostic procedures.**

---

Horton HF, Fouser L, Carter L, Ling V, Bowman MR, Carreno BM, Collins M, Wood CR, Honjo T. Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med.* 2000 Oct 2;192(7):1027-34.

Agata Y, Kawasaki A, Nishimura H, Ishida Y, Tsubata T, Yagita H, Honjo T. Expression of the PD-1 antigen on the surface of stimulated mouse T and B lymphocytes. *Int Immunol.* 1996 May;8(5):765-72.

### Related Products

12-9949 Anti-Mouse/Rat CD278 (ICOS) PE (C398.4A)

17-4031 Rat IgG2b K Isotype Control APC