

Product Data Sheet

Purified anti-mouse FR4 (Folate Receptor 4)

Catalog # / Size: 125104 / 200 µg

Clone: TH6

Isotype: Rat IgG2b, κ

Reactivity: Mouse Folate Receptor 4

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.

Applications:

Applications: FC - Quality tested

IP - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is $\leq 0.25~\mu g$ per 10^6 cells in 100 μl volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application References: 1. Yamaguchi T et al. 1007: Immunity 27(1)145

Description: Folate receptor 4 (FR4) is a surface receptor for folic acid (Vitamin B9). It was recently reported that FR4 is highly and

constitutively expressed on mouse CD4+ CD25+ natural regulatory T cells (Treg). The high expression of this surface marker on Treg can be used in combination with CD4 and CD25 to distinguish Treg from other types of T cells. Since FR4 is a surface marker, this antibody may be used in combination with anti-CD4 and/or CD25 to sort living T reg cells

for functional studies.


Antigen References: 1. Yamaguchi T et al 2007. Immunity 27:145

2. Spiegelstein O et al. 2000 Gene 258:117

Related Products: Product Clone **Application** Cell Staining Buffer

FC, ICC, ICFC FC, ICFC RBC Lysis Buffer (10X)

Purified Rat IgG2b, κ Isotype Ctrl RTK4530 FC, ICC, ICFC, IF, IHC, IP, WB



C57BL/6 mouse splenocytes stained with CD3 (145-2C11) APC and purified TH6, followed by anti-rat IgG PE.



