

## **Product Data Sheet**

## **Biotin anti-mouse FR4 (Folate Receptor 4)**

Catalog # / Size: 125105 / 50 µg

Clone: TH6

**Isotype:** Rat IgG2b, κ

Reactivity: Mouse Folate Receptor 4

Preparation: The antibody was purified by affinity chromatography, and conjugated with

biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C. **Do not freeze.** 

## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is  $\leq 0.25~\mu g$  per  $10^6$  cells in  $100~\mu l$  volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application References: 1. Yamaguchi T et al. 1007: Immunity 27(1)145

Description: Folate receptor 4 (FR4) is a surface receptor for folic acid (Vitamin B9). It was recently reported that FR4 is highly and

constitutively expressed on mouse CD4+ CD25+ natural regulatory T cells (Treg). The high expression of this surface marker on Treg can be used in combination with CD4 and CD25 to distinguish Treg from other types of T cells. Since FR4 is a surface marker, this antibody may be used in combination with anti-CD4 and/or CD25 to sort living T reg cells


for functional studies.

Antigen References: 1. Yamaguchi T et al 2007. Immunity 27:145

2. Spiegelstein O et al. 2000 Gene 258:117

**Application** Related Products: Product Clone Biotin Rat IgG2b, κ Isotype Ctrl RTK4530

FC, ICFC FC, ICFC FC, ICFC Cell Staining Buffer RBC Lysis Buffer (10X) TruStain fcX™ (anti-mouse CD16/32) 93



BALB/c mouse splenocytes stained with CD3 (145-2C11) APC and biotinylated TH6, followed by Sav-PE



